

10/530465

TITLE OF THE INVENTION

METHOD OF PREPARING INHIBITORS OF PHOSPHODIESTERASE-4

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention is directed to a method of preparing inhibitors of phosphodiesterase-4. In particular, this invention relates to a method of preparing compounds that are biaryl substituted 1,8-naphthyridin-4(1*H*)-ones.

RELATED BACKGROUND

Hormones are compounds that variously affect cellular activity. In many respects, hormones act as messengers to trigger specific cellular responses and activities. Many effects produced by hormones, however, are not caused by the singular effect of just the hormone.

Instead, the hormone first binds to a receptor, thereby triggering the release of a second compound that goes on to affect the cellular activity. In this scenario, the hormone is known as the first messenger while the second compound is called the second messenger. Cyclic adenosine monophosphate (adenosine 3', 5'-cyclic monophosphate, "cAMP" or "cyclic AMP") is known as a second messenger for hormones including epinephrine, glucagon, calcitonin, corticotrophin, lipotropin, luteinizing hormone, norepinephrine, parathyroid hormone, thyroid-stimulating hormone, and vasopressin. Thus, cAMP mediates cellular responses to hormones. Cyclic AMP also mediates cellular responses to various neurotransmitters.

Phosphodiesterases ("PDE") are a family of enzymes that metabolize 3', 5' cyclic nucleotides to 5' nucleoside monophosphates, thereby terminating cAMP second messenger activity. A particular phosphodiesterase, phosphodiesterase-4 ("PDE4", also known as "PDE-IV"), which is a high affinity, cAMP specific, type IV PDE, has generated interest as potential targets for the development of novel anti-asthmatic and anti-inflammatory compounds. PDE4 is known to exist as at least four isoenzymes, each of which is encoded by a distinct gene. Each of the four known PDE4 gene products is believed to play varying roles in allergic and/or inflammatory responses. Thus, it is believed that inhibition of PDE4, particularly the specific PDE4 isoforms that produce detrimental responses, can beneficially affect allergy and inflammation symptoms. It would be desirable to provide novel compounds and compositions that inhibit PDE4 activity.

A major concern with the use of PDE4 inhibitors is the side effect of emesis which has been observed for several candidate compounds as described in C.Burnouf et al., ("Burnouf"), *Ann. Rep. In Med. Chem.*, 33:91-109(1998). B.Hughes et al., *Br. J.Pharmacol.*, 118:1183-1191(1996); M.J.Perry et al., *Cell Biochem. Biophys.*, 29:113-132(1998);
5 S.B.Christensen et al., *J.Med. Chem.*, 41:821-835(1998); and Burnouf describe the wide variation of the severity of the undesirable side effects exhibited by various compounds. As described in M.D.Houslay et al., *Adv. In Pharmacol.*, 44:225-342(1998) and D.Spina et al., *Adv. In Pharmacol.*, 44:33-89(1998), there is great interest and research of therapeutic PDE4 inhibitors.

10 International Patent Publication WO9422852 describes quinolines as PDE4 inhibitors. International Patent Publication WO9907704 describes 1-aryl-1,8-naphthylidin-4-one derivatives as PDE4 inhibitors.


A.H.Cook, et al., *J.Chem. Soc.*, 413-417(1943) describes gamma-pyridylquinolines. Other quinoline compounds are described in Kei Manabe et al., *J.Org. Chem.*, 58(24):6692-6700(1993); Kei Manabe et al., *J.Am. Chem. Soc.*, 115(12):5324-
15 5325(1993); and Kei Manabe et al., *J.Am. Chem. Soc.*, 114(17):6940-6941(1992).

Compounds that include ringed systems are described by various investigators as effective for a variety of therapies and utilities. For example, International Patent Publication No. WO 98/25883 describes ketobenzamides as calpain inhibitors, European Patent Publication
20 No. EP 811610 and U.S. Patent Nos. 5,679,712, 5,693,672 and 5,747,541 describe substituted benzoylguanidine sodium channel blockers, U.S. Patent No. 5,736,297 describes ring systems useful as a photosensitive composition.

U.S. Patent Nos. 5,491,147, 5,608,070, 5,622,977, 5,739,144, 5,776,958, 5,780,477, 5,786,354, 5,798,373, 5,849,770, 5,859,034, 5,866,593, 5,891,896, and International
25 Patent Publication WO 95/35283 describe PDE4 inhibitors that are tri-substituted aryl or heteroaryl phenyl derivatives. U.S. Patent No. 5,580,888 describes PDE4 inhibitors that are styryl derivatives. U.S. Patent No. 5,550,137 describes PDE4 inhibitors that are phenylaminocarbonyl derivatives. U.S. Patent No. 5,340,827 describes PDE4 inhibitors that are phenylcarboxamide compounds. U.S. Patent No. 5,780,478 describes PDE4 inhibitors that are
30 tetra-substituted phenyl derivatives. International Patent Publication WO 96/00215 describes substituted oxime derivatives useful as PDE4 inhibitors. U.S. Patent No. 5,633,257 describes PDE4 inhibitors that are cyclo(alkyl and alkenyl)phenyl-alkenyl (aryl and heteroaryl) compounds.

SUMMARY OF THE INVENTION


In one aspect, the present invention is directed to a one pot method of preparing intermediates of Formula V, which are useful in making inhibitors of phosphodiesterase-4:




V

In a second aspect, the present invention is directed to a method of preparing compounds of Formula IX and Formula IXa which are inhibitors of phosphodiesterase-

4:



IX



IXa


Inhibitors of phosphodiesterase-4 have been shown to be useful in the treatment in mammals of, for example, asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal

conjunctivitis, arterial restenosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer,

5 cachexia, muscle wasting, depression, memory impairment, monopolar depression, acute and chronic neurodegenerative disorders with inflammatory components, Parkinson disease, Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour growth and cancerous invasion of normal tissues.

10 DETAILED DESCRIPTION OF THE INVENTION

In one embodiment the preparation of compounds of Formula IX may begin with the preparation of an intermediate of Formula V:



15

wherein

-OR¹ is a suitable leaving group such as those disclosed in standard texts, such as: Protective Groups in Organic Synthesis, 2nd edition, by Theodora W. Greene and Peter G M. Wuts (1991) and Protecting Groups, by Philip J. Kocienski (1994). For example, R¹ may be an optionally C₁-8alkyl, aryl, heteroaryl, wherein the substituents are selected from the group consisting of aryl, C₁-8 alkyl;


20

R³ is C₁-8alkyl, aryl or heteroaryl, optionally mono, di-, tri- or tetra-substituted with Halo, B(OH)₂, —C≡CH, aryl, heteroaryl, C₁-10alkyl, alkoxy, ketone, alcohol, triflate, mesylate, amide, ester, ketone, acetal, phosphate, stannane, silyl, B(OR)₂ wherein R is C₁-

25 galkyl or aryl;

comprising

Step A: reacting, in a non-nucleophilic solvent, a compound of Formula II:




II

5

wherein

R² is each independently C₁-alkyl, optionally mono or di-substituted with C₁-alkyl, such as methyl or ethyl, or aryl, such as phenyl or heteroaryl;

10 with a compound of Formula I in the presence of a base:




I

wherein

15 R⁴ is bromo or chloro,


to yield a compound of Formula III.



III

5 and,

Step B: reacting, in a non-nucleophilic solvent, a compound of Formula III with an amine compound of Formula IV



IV

10

in the presence of a base to yield a compound of Formula V.

The definition for the non-nucleophilic solvent of Step B is as previously defined for Step A. Similarly, the definition for the base of Step B is as previously defined for Step A.

15

In Step A, the molar ratio of base to compound of Formula I can be varied from about 12:1 to about 3:1. A ratio of 8:1 is typical. A ratio lower than 3:1 may result in trimerization of the aminoacrylate to give 1,3,5-tricarboxybenzene. The molar ratio of the compound of Formula I to compound of Formula II can be varied from 1:1 to 1:3; typically about 1:1.5. Reaction Step A may conveniently be conducted at a temperature range of 25 to 100°C; or typically 40 to about 60°C and is allowed to proceed until substantially complete in from 2 to 18 hours; typically 6 to 12 hours. Reaction Step B is carried out as described above.

20

Within this embodiment there is a genus wherein:

R^1 is an optionally C_{1-8} alkyl, aryl, heteroaryl, wherein the substituents are selected from the group consisting of aryl, C_{1-8} alkyl;

25

R^2 is methyl or ethyl or phenyl;

R^3 is C_{1-8} alkyl, aryl or heteroaryl, optionally mono, di-, tri- or tetra-substituted with Halo, $B(OH)_2$, $-C\equiv CH$, aryl, heteroaryl;

Within this embodiment there is a genus wherein:

R¹ is methyl or ethyl;

R² is methyl;

R³ phenyl substituted with halo, B(OH)₂, —C≡CH.

Within this embodiment there is a genus wherein:

the non-nucleophilic solvent is selected from dimethylformamide, dimethylacetamide, tetrahydrofuran, ethylacetate, acetonitrile, toluene, benzene, dioxane, methylene chloride

Within the genus immediately above there is a sub-genus wherein:

the non-nucleophilic solvent is acetonitrile.

Within this embodiment there is a genus sub-genus wherein:

the base is selected from sodium or potassium carbonate, triethylamine, diethylamine, and Huenigs base.

Within this embodiment there is a genus wherein:

the base is selected from sodium carbonate or triethylamine.

Within this embodiment there is a genus wherein:

the molar ratio of compound of Formula IV to compound of Formula III is about 2:1 to 1:2.

Within the genus immediately above there is a sub-genus wherein:

the molar ratio of the compound of Formula IV to the compound of Formula III is about 1:1.

Within this embodiment there is a genus wherein:

the molar ratio of base to compound of Formula IV is at least 1:1.

Within the genus immediately above there is a sub-genus wherein:

the molar ratio of base to compound of Formula IV is about 1:1.


Within this embodiment there is a genus wherein:

the reaction is conducted at a temperature range of about 20 to 100°C.

Within the genus immediately above there is a sub-genus wherein:

the reaction is conducted at a temperature range of about 40 to about 50°C.

In a second embodiment there is a method of preparing a compound of Formula V



V

5 wherein


-OR¹ is a suitable leaving group;

R³ is C₁-8alkyl, aryl or heteroaryl, optionally mono, di-, tri- or tetra-substituted with Halo, B(OH)₂, —C≡CH (alkyne), aryl, heteroaryl, C₁-10alkyl, alkoxy, ketone, alcohol, triflate, mesylate, amide, ester, ketone, acetal, phosphate, stannane, silyl, B(OR)₂ wherein R is

10 C₁-8alkyl or aryl;

comprising

Step A: reacting, in a non-nucleophilic solvent, a compound of Formula II:



II


15

wherein

R² is each independently C₁-8alkyl, optionally mono or di-substituted with C₁-8alkyl, such as methyl or ethyl, or aryl or heteroaryl;

20

with a compound of Formula I in the presence of a base:




I

wherein

R⁴ is bromo or chloro

5

to yield a compound of Formula III.



III

10 and,

Step B: reacting, in a non-nucleophilic solvent, a compound of Formula III with an amine compound of Formula IV



IV

15

in the presence of a base to yield a compound of Formula V.

Within this second embodiment there is a genus wherein:

20 R¹ is an optionally C₁-8alkyl, aryl, heteroaryl, wherein the substituents are selected from the group consisting of aryl, C₁-8 alkyl;

R² is methyl or ethyl or phenyl;

R³ is C₁-8alkyl, aryl or heteroaryl, optionally mono, di-, tri- or tetra-substituted with Halo, B(OH)₂, —C≡CH, aryl, heteroaryl;

Within this second embodiment there is a genus wherein:

R⁴ is chloro.

Within this second embodiment there is a genus wherein:

R¹ is methyl or ethyl;

R² is methyl;

R³ phenyl substituted with halo, B(OH)₂, —C≡CH.

Within this second embodiment there is a genus wherein:

the non-nucleophilic solvent is selected from dimethylformamide, dimethylacetamide, tetrahydrofuran, ethylacetate, acetonitrile, toluene, benzene, dioxane, methylene chloride

Within this genus there is a sub-genus wherein:

the non-nucleophilic solvent is acetonitrile.

Within this second embodiment there is a genus wherein:

the base is selected from sodium or potassium carbonate, triethylamine, diethylamine, and Huenigs base.

Within this genus there is a sub-genus wherein:

the base is selected from sodium carbonate or triethylamine.

Within this second embodiment there is a genus of Step A, wherein:

the molar ratio of base to compound of Formula I is about 12:1 to about 3:1.

Within this genus, there is a sub-genus of Step A, wherein:

the molar ratio of base to compound of Formula I is about 8:1 to 3:1.

Within this second embodiment, there is a genus of Step A, wherein:

the molar ratio of the compound of Formula I to compound of Formula II is about 1:1 to 1:3.

Within this second embodiment, there is a genus of Step A, wherein:

the reaction is carried out at about 25 to 100°C.

Within this second embodiment, there is a genus Step A wherein:

The reaction is carried out at about 40 to about 60°C.

Within this second embodiment, there is a genus of Step B, wherein:

the molar ratio of compound of Formula IV to compound of Formula III is about 2:1 to 1:2.

Within this genus there is a sub-genus of Step B wherein:

the molar ratio of the compound of Formula IV to the compound of Formula III is about 1:1.

Within this second embodiment there is a genus of Step B, wherein:

the molar ratio of base to compound of Formula IV is about 1:1 or more.

Within this genus, there is a sub-genus of Step B, wherein:



the molar ratio of base to compound of Formula IV is about 1:1.

Within this second embodiment there is a genus of Step B, wherein:
the reaction is conducted at a temperature range of about 20 to 100°C.

- 5 Within this genus there is a sub-genus of Step B, wherein:
the reaction is conducted at a temperature range of about 40 to about 50°C.


Within this second embodiment, there is a genus wherein:
reaction step A and Reaction Step B are carried out in a single pot without purification or
isolation of the product of Step A prior to proceeding with Step B.

- 10 In another embodiment, the present invention is direct to a method of preparing a
phosphodiesterase-4 inhibitor of Formula IX and IXa



- 15 Comprising


Step C: reacting, in solvent A, a compound of Formula Va




wherein

-OR¹ is a suitable leaving group such as those disclosed in standard texts, such as: Protective Groups in Organic Synthesis, 2nd edition, by Theodora W. Greene and Peter G M. Wuts (1991) and Protecting Groups, by Philip J. Kocienski (1994). For example, R¹ may be an optionally C₁-8 alkyl, aryl, heteroaryl, wherein the substituents are selected from the group consisting of aryl, C₁-8 alkyl;

with a compound of Formula VII or VIIa




VII




VIIa

in the presence of a palladium catalyst and a phosphine ligand in the amine base to yield a compound of Formula VIII or VIIIa




VIII



VIIIa

A preferred compound of Formula Va and Formula VIII and VIIIa, respectively, are as follows:



For purposes of this specification, solvent A is selected from the group consisting of dimethylaminoacetamide, dimethylformamide, acetonitrile, DMSO, methylacetamide, ethers or mixtures thereof. For purposes of this specification, the phosphine ligand is selected from the group consisting of $P(C_1-6alkyl)_3$, such as $P(t-butyl)_3$, $P(Cy)_3$, and $P(t-butyl)_2(biphenyl)$. For purposes of this specification, the palladium catalyst includes Fu's catalyst (i.e. $P(t-butyl)_3Pd-P(t-butyl)_3$), $[PdCl(allyl)]_2$, $Pd_2(dba)_3$, and $[P(t-butyl)_3PdBr]_2$ (Johnson-Matthey catalyst). For purposes of this specification, the amine base is a hindered amine base such as t-amylamine, t-butylamine, isopropylamine, or diisopropylamine.

In Step C, the molar ratio of the compound of Formula Va to Formula VII or VIIa may vary widely, but is optimally about 1:1.5 to 1.5:1 and is typically approximately 1:1. As a general principle, at least 1 molar equivalent of base per mole of Compound of Formula Va plus the 1 molar equivalent of base per molar equivalent of the compound of Formula VII or VIIa is desirable. Reducing this ratio reduces the yield. Thus, in the reaction above the ratio of amine base to compound of Formula Va can be 1.2:1 or greater, but is typically 2:1 to 3.5:1. The ratio molar of Palladium to compound of Formula V 0.05:1 to 0.10:1. In the instance $[PdCl(allyl)]_2$, for example, which possesses two mole of palladium per mole of catalyst, the ratio would be 0.025:1 to 0.05:1. The molar ratio of phosphine ligand to palladium is typically 3:1 to 5:1, often approximately 4:1. The reaction can be carried out a 25 to 125°C, more typically 40-70°C, and is allowed to proceed until the reaction is substantially complete. The presence of oxygen is minimized by purging the reaction vessel with nitrogen gas prior to addition of the reagents.

The product of Step C is cooled to 0 to about 5°C. Thereafter 3-6 volumes of anti-solvent per volume of solvent is added to the product of Step C to precipitate compound of

Formula VIII or VIIIa. For purposes of this specification, the anti-solvent is defined as a non-reactive solvent in which compound of Formula VIII or VIIIa will precipitate.

The slurry comprising the precipitated compound of Formula VIII or VIIIa is filtered or centrifuged. The filter or centrifuge cake comprising the compound of Formula VIII or VIIIa is re-slurried or directly washed with isobutanol or isopropanol to remove residual palladium. For example, the filter cake comprising compound of Formula VIII or VIIIa may be washed in situ (i.e. on the filter press) with 40 to 60 volumes of isobutanol unit weight of dry compound of Formula VIII or VIIIa.

As an alternative, residual palladium may be removed by re-dissolving compound of Formula I in an C₁₋₆alkanol solvent (such as methanol, ethanol or propanol) and subjecting the slurry comprising the compound of Formula IX or IXa to an activated carbon such as Darco G-60 or Ecosorb. The slurry is then filtered to remove the activated carbon. As may be appreciated, it is convenient to increase the solubility of the product in the alkanol solvent by warming the alkanol (such as to 25 to 70°C) either before or after contact with the product. Typically, the slurry is kept warm for 1 to 6 hours, and filtered while still warm.

In function, the overall transformation achieved in step C is the same as that of a typical Sonogashira coupling reaction. As such, Step C is an efficient, one step method of coupling an alkyne to an aryl halide. On the other hand, step C differs from a Sonogashira coupling reaction in several important ways. For example, the Sonogashira coupling reaction typically requires the use of a copper, zinc, magnesium or zirconium reagent in addition to a palladium catalyst. Copper is preferred. However, applicants have found that Sonogashira coupling (Sonogashira, K. in *Metal-Catalyzed Cross-Coupling Reactions*; Diederich, F. and Stang, P. eds; Wiley-VCH: Weinheim, Germany, 1998, Chapter 5. See also, Littke, A. and Fu, G. *Angew. Chem. Int. Ed.* 1998, 37, No. 24, pp 3387-3388; Wolfe, J. Singer, R., Yang, B and Buchwald, S., *J. Am. Chem. Soc.*, 1999, 121, pp 9550-9561.) of compound of Formula Va and Formula VII results of the production of as much as 10% dimer of the compound of Formula Va in the product.

Hermann, et. al. (*Eur. J. Chem* 2000, 3679-3681), reported that if the palladium catalyst is judiciously chosen, the need for copper can be obviated. P(Cy)₃-Pd-P(Cy)₃ and P(t-butyl)₃-Pd-P(t-butyl)₃ (Fu's catalyst), are two of such catalysts. Such catalysts require special handling. In contrast, the catalysts of the present invention are prepared in situ upon reaction of a stable palladium source (such as [Pd(allyl)Cl]₂) with a stable commercially available, pre-packaged ligand (such a 10% by weight tri-(t-butyl)phosphine in hexane. Moreover, by the

creation of the catalyst, in situ, the invention provides convenient access to ligand/Palladium ratio's generally unavailable through commercial sources.

Step D: reacting, in solvent B a compound of Formula VIII with
5 cyclopropylamine, optionally in the presence of a phosphite or other catalyst.

For purposes of this specification solvent B is defined to include an alkanol solvent such C₁₋₄alkanol solvents methanol, ethanol, propanols and butanols, and trifluoroethanol, or mixtures thereof. Acetonitrile may also be used. For purposes of this
10 specification, the catalyst is defined to include Butyl phosphite (BuO)₃P, pyridine-*N*-oxides, tri-alkyl orthoacetate, zinc chloride, magnesium chloride, magnesium triflate, scandium triflate, lanthanum triflate, ytterbium triflate, titanocene dichloride, or zirconocene dichloride. Note that for purposes of this specification, the term catalyst is intended to indicate a reagent that accelerates the reaction, even if present in greater than classical "catalytic" amounts.

15 The molar ratio of cyclopropylamine to compound of Formula VIII or VIIIa is typically greater than 1:1, and if often 1.1:1 or greater. Reducing this ratio reduces the yield. The ratio of catalyst to compound of Formula VIII or VIIIa is generally 0.025:1 to 0.075: 1, often about 0.05:1 or possibly 0.10:1 to 1:1 or more. In the case of magnesium chloride a typical ratio of catalyst to compound of Formula VIII or VIIIa is 0.9:1. to 1:1.5. The reaction can
20 be carried out at 40 to about 60°C, more typically 55-60°C, and is allowed to proceed until the reaction is substantially complete. The presence of oxygen is minimized by purging the reaction vessel with nitrogen gas prior to addition of the reagents. For optimal results, it desirable to perform Step D at or below a Kf of 3000ppm.

25 The product of Step D is cooled to 0 to about 30°C, typically about room temperature. Thereafter, the solvent is replaced with an anti-solvent (anti-solvent B), such as by evaporation of the solvent, follow by the addition of excess anti-solvent. For purposes of this specification, anti-solvent B is defined as a solvent in product of the reaction (in this case the compound of Formula IX) 3-6 volumes of anti-solvent per volume of solvent is added to the product of Step C to precipitate compound of Formula VIII or VIIIa . For purposes of this
30 specification, the anti-solvent is defined as a non-reactive solvent in which compound of Formula VIII or VIIIa will precipitate.

In order to remove impurities, the slurry comprising the precipitated compound of Formula VIII or VIIIa is washed a solvent, such as water, acetonitrile, isopropyl acetate, ethyl acetate, tetrahydrofuran, ethanol, propanols, and butanols. Thereafter, the solvent is removed,

such as by filtration and vacuum drying. The filtered compound of Formula VIII or VIIIa can then be mixed with a conversion solvent, such as dry ethanol, methanol, trifluoroethanol, *N*-methylpyrrolidinone, methyl *t*-butyl ether or mixtures thereof, optionally heated to and held at 40 to 50°C until at least 95% of the compound of Formula VIII or VIIIa is in the desired form. The conversion solvent is then removed, such as by filtration or centrifugation and the resulting product is vacuum dried. The amount of conversion solvent required, is that sufficient to suspend or dissolve the compound of Formula VIII or VIIIa.

The term "aryl" unless specifically stated otherwise includes multiple ring systems as well as single ring systems such as, for example, phenyl or naphthyl.

The term "hetero" unless specifically stated otherwise includes one or more O, S, or N atoms. For example, heterocycloalkyl and heteroaryl include ring systems that contain one or more O, S, or N atoms in the ring, including mixtures of such atoms. The hetero atoms replace ring carbon atoms. Thus, for example, a heterocycloC₅alkyl is a five membered ring containing from 5 to no carbon atoms.

Examples of heteroaryl include, for example, pyridinyl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinoxalinyl, furyl, benzofuryl, dibenzofuryl, thienyl, benzothienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl.

Compounds described herein can contain one or more asymmetric centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and salts thereof.

The utility of the invention is illustrated by examples of phosphodiesterase-4 inhibitors that may be prepared by use of the invention.

Dosage levels from about 0.001mg/kg to about 140mg/kg of body weight per day are useful in the treatment of conditions such as asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus

host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, monopolar depression, acute and chronic neurodegenerative disorders with inflammatory components, Parkinson disease, Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour growth and cancerous invasion of normal tissues which are responsive to PDE4 inhibition, or alternatively about 0.05mg to about 7g per patient per day. For example, inflammation may be effectively treated by the administration of from about 0.01mg to 50mg of the compound per kilogram of body weight per day, or alternatively about 0.5mg to about 2.5g per patient per day. Further, it is understood that the PDE4 inhibiting compounds of this invention can be administered at prophylactically effective dosage levels to prevent the above-recited conditions.


The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration to humans may conveniently contain from about 0.5mg to about 5g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms will generally contain between from about 0.01mg to about 1000mg of the active ingredient, typically 0.01mg, 0.05mg, 0.25mg, 1mg, 5mg, 25mg, 50mg, 100mg, 200mg, 300mg, 400mg, 500mg, 600mg, 800mg or 1000mg.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The examples that follow are intended as an illustration of certain preferred embodiments of the invention and no limitation of the invention is implied.


Compounds of the present invention can be prepared according to the following methods. The substituents are the same as in Formula I except where defined otherwise.

Preparation of Intermediate compound of Formula V.



5

NAPHTHYRIDONE 1




Step 1: Ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate.

A mixture of ethyl 2-chloronicotinoyl acetate (purchased or prepared following a procedure described in J. Het. Chem. ,30, 855, 1993) (1 eq), triethylamine (4 eq) and ethyl 3,3-dimethylaminoacrylate (1.5 eq) in acetonitrile (0.5M) was heated to reflux for 3h, cooled to 40-50°C and 3-bromoaniline (1 eq) was added. The reaction was heated to reflux overnight, cooled to rt, diluted with water (2 volume). The product was isolated by filtration and washed with water, ether or acetonitrile-water (1:1).

¹H NMR (Acetone-d₆) δ 1.32 (t, 3H), 4.29 (q, 2H), 7.54-7.63 (m, 2H), 7.69 (dd, 1H), 7.78 (dd, 1H), 7.93 (s, 1H), 8.66-8.71 (m, 3H).

NAPHTHYRIDONE 2




Ethyl-1-(3-acetylenylphenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 3-acetylenylaniline for 3-bromoaniline results in the Title Naphthyridone 2.

¹H NMR (CDCl₃) δ 1.42 (t, 3H), 3.19 (s, 1H), 4.20 (q, 2H), 7.42-7.46 (m, 2H), 7.53-7.59 (m, 2H), 7.67 (dt, 1H), 8.64 (dd, 1h), 8.68 (s, 1H), 8.83 (dd, 1H).


NAPHTHYRIDONE 3

5 Ethyl-1-(3-cyanophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 3-cyanoaniline for 3-bromoaniline results in the Title Naphthyridone 3.

10 ^1H NMR (DMSO- d_6) δ 1.27 (t, 3H), 4.23 (q, 2H), 7.57 (dd, 1H), 7.80 (t, 1H), 7.96-8.04 (m, 2H), 8.20 (t, 1H), 8.61 (dd, 1H), 8.68 (dd, 1H), 8.70 (s, 1H).

NAPHTHYRIDONE 4




15

Ethyl-1-(4-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 4-bromoaniline for 3-bromoaniline results in the Title Naphthyridone 4.

20 ^1H NMR (DMSO- d_6) δ 1.26 (t, 3H), 4.22 (q, 2H), 7.54-7.59 (m, 3H), 7.78 (d, 2H), 8.61 (dd, 1H), 8.63 (s, 1H), 8.68 (dd, 1H).


NAPHTHYRIDONE 5

5 Ethyl-1-(2-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 2-bromoaniline for 3-bromoaniline results in the Title Naphthyridone 5.

10 ^1H NMR (DMSO- d_6) δ 1.26 (t, 3H), 4.23 (q, 2H), 7.53-7.62 (m, 3H), 7.75 (dd, 1H), 7.88 (dd, 1H), 8.61 (s, 1H), 8.63 (dd, 1H), 8.68 (dd, 1H).

NAPHTHYRIDONE 6



15


Ethyl-1-(2-*tert*-butylphenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 1-bromo-2-*tert*-butylphenylaniline for 1-bromo-2-*tert*-butylaniline results in the Title Naphthyridone 6.

20

^1H NMR (DMSO- d_6) δ 1.08 (s, 9H), 1.25 (t, 3H), 4.22 (q, 2H), 7.30-7.40 (m, 2H), 7.50-7.56 (m, 2H), 7.73 (dd, 1H), 8.59 (s, 1H), 8.62 (dd, 1H), 8.70 (dd, 1H).

NAPHTHYRIDONE 7

5 Ethyl-1-(3-phenylboronic acid)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure described for Naphthyridone 1, but substituting 3-aminophenylboronic acid for 3-bromoaniline results in the Title Naphthyridone 7.

10 ^1H NMR (DMSO- d_6) δ 1.26 (t, 3H), 4.22 (q, 2H), 7.56 (coincident triplets, 2H), 7.60 (d, 1H), 7.87 (s, 1H), 7.94 (d, 1H), 8.24 (s, 2H, exchangeable), 8.61 (d, 1H), 8.62 (s, 1H), 8.68 (d, 1H).

PDE-4 INHIBITOR COMPOUNDS

15 The following are examples of PDE-4 inhibitor compounds which can be made following the procedures as described herein. Note that the formula number used in the schemes 1 to 12 and the discussion of these schemes should not be confused with the formula numbers used elsewhere in this patent application. Examples 1 to 13 were prepared using Naphthyridone 1:

20

Scheme 1

25 In a first method outlined in Scheme 1 below, an appropriately substituted derivative of ethyl 2-chloronicotinoyl acetate of formula II is reacted with 1.5 equivalents of triethyl orthoformate and 5 equivalents of acetic anhydride at 130°C, and after removal of the volatile components, the crude 2-chloronicotinoyl acrylate of formula III is immediately reacted with 1.2 equivalents of an appropriately substituted haloaryl amine of formula IV, such as, for example 3-bromoaniline, in a halogenated hydrocarbon solvent such as methylene chloride at a temperature of 0°C to room temperature. After an appropriate reaction time ranging from 2 to 24

30 hours the resulting 3-arylamino acrylate of formula V is obtained by evaporation of the solvent

and may be further purified by chromatography on silica gel or crystallization from an appropriate solvent.

The compound of formula V may alternatively be used without further purification in the following step. Cyclization of the compound of formula V to the 1-haloaryl-1,4-dihydro[1,8]naphthyridin-4-one carboxylate of formula VI is effected by treatment with a small excess of a strong base such as an alkali metal hydride, for example sodium hydride, in an appropriate solvent such as tetrahydrofuran at a starting temperature of 0°C with warming to room temperature if required to complete the process. The product of formula VI is isolated in crude form by dilution with a large volume of water followed by filtration or by extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

The product of formula VI thus obtained can be hydrolyzed to the corresponding carboxylic acid derivative under basic conditions, using an aqueous solution of an alkali base such as an alkali carbonate or preferably sodium or potassium hydroxide, with an organic cosolvent such as tetrahydrofuran or a primary, secondary or tertiary alkanol, such as methanol or ethanol, or a combination thereof at temperatures ranging from room temperature to reflux temperature for the appropriate time. The resultant carboxylic acid is isolated in crude form following acidification using an aqueous solution of an inorganic acid such as hydrochloric, sulfuric or a similar acid, and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

The carboxylic acid is then transformed into the appropriate primary, secondary or tertiary amide analog of formula VII by any general procedure well known to the organic chemist, preferably via initial transformation into a mixed anhydride by treatment with a small excess, such as 1.25 equivalents, of an appropriate alkyl chloroformate such as ethyl or isobutyl chloroformate, in the presence of a larger excess, such as 2.5 equivalents, of a tertiary organic amine such as triethylamine or N,N-diisopropylethylamine in an organic solvent such as tetrahydrofuran at low temperature, preferably 0°C, for a period of 30 minutes to 3 hours. An excess, usually 5 or more equivalents, of an appropriate primary or secondary amine or of an aqueous solution of ammonium hydroxide is then added and the resulting reaction is allowed to

proceed at a temperature ranging from 0°C to room temperature for an appropriate length of time, usually 1-24 hours.

The desired amide of formula VII is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration. In cases where the amide moiety is 2,6-dichloropyridin-4-yl, a different procedure is used in which the anion of 4-amino-3,5-dichloropyridine is generated at low temperature, preferably at 0°C, using a strong alkali hydride such as sodium hydride in a solvent such as tetrahydrofuran, and reacted with the acid chloride of a carboxylic acid (from hydrolysis of an ester of formula VI) generated by an appropriate known procedure, usually by the action of oxalyl chloride activated by a catalytic amount of N,N-dimethylformamide in a solvent such as tetrahydrofuran.

The amides of general formula VII are processed into the products of formula I by reaction with an appropriately substituted aryl or heteroaryl boronic acid or boronate ester of formula VIII under the catalysis of a transition metal species such as *trans*-dibromobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in an appropriate solvent or solvent mixture, preferably a 1:1 mixture of toluene and ethanol in the presence of an excess of an aqueous solution of an alkali base such as sodium carbonate, at an appropriate temperature, preferably 50 to 100°C, for an appropriate length of time ranging from 0.5 to 48 hours.


The resulting reaction product is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

Compounds of formula I may also be obtained by reaction of a compound of formula VII with an appropriately substituted aryl or heteroaryl tributyl stannane of formula IX under the catalysis of a transition metal species such as *trans*-dibromobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in the presence of a copper (I) species such as cuprous iodide in an appropriate solvent such as N,N-dimethylformamide at a temperature range of 50-100°C for a period of 2 to 24 hours. Isolation of the reaction product is effected as described above.

Alternatively, an ester of formula VI can be processed into an ester of formula X by reaction with an appropriately substituted boronic acid or boronate ester, or with an appropriately substituted stannane derivative under the conditions described above, and the ester can be hydrolyzed and transformed into an amide of formula I.

- 5 The boronic acids of formula VIII or corresponding boronate esters are usually obtained from commercial sources. Where required, they can be prepared readily from the corresponding halides via metallation with n-butyllithium followed by reaction with a trialkyl borate, or by using classical transition metal-catalyzed coupling procedures using diboron pinacol ester. The stannanes of formula IX are generated from the corresponding halides via
10 initial metallation using n-butyllithium followed by addition of tributyltin chloride.

Scheme 1




In an alternative method for the preparation of compounds of formula I, outlined in Scheme 2 below, an amide of formula VII can be transformed into a corresponding boronate ester of formula XI by treatment with an excess of diboron pinacol ester in the presence of an inorganic salt such as potassium acetate under the catalysis of a transition metal species such as *trans*-dibromobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in a solvent such as N,N-dimethylformamide at temperatures ranging from 50 to 100°C for a length of time ranging from 1 to 48 hours. The boronate of formula XI may be isolated by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The resulting product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

Alternatively, the boronate of formula XI can be used as generated *in situ* in the reaction medium without isolation, and reacted with a small excess of an appropriately substituted aryl or heteroaryl halide of formula XII under the catalysis of a transition metal species such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in an appropriate solvent or solvent mixture, preferably a 1:1 mixture of toluene and ethanol in the presence of an excess of an aqueous solution of an alkali base such as sodium carbonate, at an appropriate temperature, preferably 50 to 100°C for an appropriate length of time ranging from 0.5 to 48 hours.


The reaction product of formula I is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

Scheme 2




In a third method used for the synthesis of compounds of formula I of this invention (scheme 3), an intermediate nicotinoyl acrylate of formula III is reacted with an appropriately constructed diaryl or heteroarylaryl amine of formula XIII under the conditions described previously to afford a compound of formula XIV which is cyclized by the action of a strong base such as sodium hydride as described above to afford an ester of formula X which is processed into a compound of formula I via hydrolysis and amide formation as described above.


Scheme 3




The diaryl or heteroarylamine intermediates of formula XIII were assembled as indicated in Scheme 4. An appropriately substituted aniline boronic acid of formula XV is coupled with an appropriately substituted aryl or heteroaryl halide of general formula XII under the catalysis of a transition metal species as described above to afford the formula XIII compounds used in Scheme 3.

Scheme 4


Bromopyridine intermediates substituted at the 2-position by carbon based nucleophiles of formula XVII, where R^8 is selected from R^2 moieties having a carbon-carbon link to the pyridine, are conveniently prepared as shown in Scheme 5. The bromopyridine intermediates are prepared from dihalides of formula XVI by treatment with an appropriate solution of a Grignard reagent under the catalysis of a transition metal species such as [1,1'-bis(diphenylphosphino)ferrocene]dichloronickel (II) in a solvent such as tetrahydrofuran at a temperature range of -10°C to room temperature and the resulting reaction mixture worked up by well known procedures to afford the desired product.

Scheme 5

Halopyridine intermediates of type XVIII where the 2-substituent is an alkoxy group OR^9 are derived from dihalides of formula XVI by displacement with an appropriate alkali alkoxide as outlined in scheme 6. The reaction is effected in a solvent such as N,N-dimethylformamide at a temperature range of 0°C to room temperature and, upon completion of the reaction, the products are isolated and purified following classical procedures.


Scheme 6

- Where intermediates of formula XIX or XX in which the 2-substituent is sulfide, sulfoxide or sulfone were required, they were attained as described in Scheme 7. An appropriate dihalopyridine of type XVI is reacted with an appropriate thioalkoxide, usually generated from the corresponding thiol or disulfide through the action of a strong base such as an alkali hydride or n-butyllithium, in a solvent such as N,N-dimethylformamide or diethyl ether at a temperature range of -78°C to room temperature. Upon completion of the reaction the products of formula XIX are isolated and purified following classical procedures. The products thus obtained can be oxidized to the corresponding sulfoxides or sulfones of formula XX through the action of an oxidizing agent such as oxone or an organic peracid. In Scheme 7, R¹⁰ is H or C₁₋₆alkyl.


Scheme 7

The preparation of haloacyl pyridine intermediates of formula XXII, presented in Scheme 8, requires treatment of a halopyridine ester of type XXI with a solution of an appropriate Grignard reagent in a solvent such as diethyl ether at a temperature range of 0°C to room temperature. If the reaction is carried out for a longer period or under reflux a halopyridine carbinol of formula XXIII is obtained. In Schemes 8 and 9, R⁷ is C₁₋₆alkyl and R⁶ is methyl or ethyl.

Scheme 8




Scheme 9 outlines an alternative sequence for the synthesis of certain halopyridine carbinols of type XXIII. When 2,5-dibromopyridine is treated with n-butyllithium in toluene at -78°C followed by addition of an appropriate ketone or aldehyde and subsequent quenching at -78°C, a carbinol of type XXIII results where the carbinol group occupies the 2-position of the pyridine ring. If the metallation step is performed in diethyl ether, the same process leads to an intermediate of formula XXIII in which the carbinol group occupies the 5-position of the pyridine ring.

Scheme 9

R^7 and R^8 can be the same or different
either R^7 or R^8 can be hydrogen


Scheme 10 demonstrates the methods of synthesis for compounds of formula I in which R^2 is a substituted phenyl or heteroaryl group. An intermediate compound of Type I where R^2 is a halogen is reacted with an appropriately substituted boronic acid or boronate ester of formula VII or tributyl stannane of formula IX using one of the methods described above to afford the desired compound.

Scheme 10

Various further transformations on pre-assembled compounds of formula I are illustrated in Scheme 11. In cases where the Ar group is a pyridine or quinoline group it can be oxidized to the corresponding nitrogen oxide by the action of an appropriate oxidizing agent such as m-chloroperoxybenzoic acid or magnesium monoperoxyphthalate under commonly used conditions. In cases where one or more of the substituents on the Ar group is a ketone it is conveniently transformed into an oxime analog through the action of hydroxylamine in pyridine as solvent. A sulfide substituent is easily oxidized to the corresponding sulfoxide or sulfone derivative by using an appropriate quantity of an oxidant such as oxone or an organic peracid.


The transformation of a 2-benzyloxypyridine into the corresponding 2-pyridone was accomplished by treatment with trifluoroacetic acid in a solvent such as methylene chloride at room temperature or under slight warming. The removal of a tert-butyloxycarbonyl protecting group from a piperazine ring is effected by reaction with trifluoroacetic acid in a solvent such as 1,2-dichloroethane at reflux temperature. In examples where a substituent on Ar is a hydroxymethyl group it can be derivatized to the analogous halomethyl moiety using a tetrahalomethane in the presence of a trisubstituted phosphine such as triphenylphosphine or diphos in a solvent such as methylene chloride. The halide can be displaced by an appropriate sulfinic acid sodium salt to afford the alkyl or arylsulfonylmethyl analog.

Scheme 11



The transformation of a 1-hydroxy-1-methylalkyl derivative such as exemplified by compounds of type XXIV of scheme 12 into 1,2-dihydroxyalkyl analogs of type XXVI is effected via initial acid-catalyzed dehydration, for example by heating in aqueous sulfuric acid, to afford an intermediate 1-alkylvinyl species of type XXV which is transformed into the desired diol XXVI by a dihydroxylation process, using for example an oxidant such as 4-methylmorpholine N-oxide (NMO) in the presence of a catalytic quantity of potassium osmate dihydrate.


Scheme 12



EXAMPLES

5 EXAMPLES of the present invention are summarized in the following table referring to Formula (I):

Table 1



10

EX.	R	R ¹	R ²	R ³	R ⁴	R ⁶
1	H	i-pr	3-C(O)Me	H	H	H
2	H		3-C(O)Me	H	H	H
3	H	i-pr	4-n-propyl	H	H	H
4	H	i-pr	4-C(O)Me	H	H	H




EX.	R	R ¹	R ²	R ³	R ⁴	R ⁶
5	H	i-pr	2-Me	H	H	H
6	Me	i-pr	4-C(O)Me	H	H	H
9	H	t-bu	4-C(O)Me	H	H	H
11	H	i-pr	4- 	H	H	H
16	H	c-pr	4-CH ₂ OH	H	H	H
18	H	c-pr	4-SEt	H	H	H
20	H	c-pr	4-SO ₂ NH ₂	H	H	H
21	H	i-pr	3-OEt	H	H	H
22	H	i-pr	4-SMe	H	H	H
23	H	i-pr	3-C(O)Me	4-OH	H	H
49	H	i-pr	4-SO ₂ Me	H	H	H
52	H	c-pr	4-SO ₂ Et	H	H	H
53	H	c-pr	4-S(O)Et	H	H	H
54	H	i-pr	4-C(=NOH) Me	H	H	H
55	H	i-pr	4- 	H	H	H
56	H	c-pr	4-CH ₂ SO ₂ Me	H	H	H

Table 2



EX.	R	R ¹	π - Position	R ²	R ³	R ⁴	R ⁶	n
7	H	i-pr	3	H	H	H	H	0
10	H		3	H	H	H	H	0
14	H	c-pr	3	H	H	H	H	0
15	H	i-pr	3	5-SMe	H	H	H	0
17	H	c-pr	4	H	H	H	H	0
24	H	i-pr	3	5-COOEt	H	H	H	0
25	H	i-pr	3	5-CMe ₂ OH	H	H	H	0
26	H	i-pr	3	6-CH ₂ CHMe ₂	H	H	H	0
27	H	i-pr	3	5-C(O)Me	H	H	H	0
28	H	i-pr	3	6-Me	H	H	H	0
30	H	H	3	6-CMe ₂ OH	H	H	H	1
32	H	c-pr	3	5-SO ₂ Me	H	H	H	0
33	H	c-pr	2	4-CMe ₂ OH	H	H	H	1
34	H	c-pr	2	5-CMe ₂ OH	H	H	H	0
35	H	c-pr	4	3-CMe ₂ OH	H	H	H	0
36	H	c-pr	4	3-CMe ₂ OH	H	H	H	1



EX.	R	R ¹	π - Position	R ²	R ³	R ⁴	R ⁶	n
37	H	c-pr	3	6-SO ₂ I-pr	H	H	H	0
38	H	c-pr	3	6-OMe	H	H	H	0
39	H	c-pr	3	6-Me	H	H	H	0
40	H	c-pr	3	6-OCH ₂ CF ₃	H	H	H	0
41	H	c-pr	3	5-Br	H	H	H	0
42	H	c-pr	3	6-OCH ₂ Ph	H	H	H	0
43	H	c-pr	3	6-C(c-pr) ₂ OH	H	H	H	0
44	H	c-pr	2	5-CMe ₂ OH	H	H	H	1
45	H	c-pr	3	6-CMe ₂ OH	H	H	H	0
46	H	i-butyl	3	6-CMe ₂ OH	H	H	H	0
47	H	c-pr	3	6-CMe ₂ OH	H	H	5-Br	0
48	H	c-pr	2	6-CMe ₂ OH	H	H	H	0
50	H	c-pr	3	6-SO ₂ Me	H	H	H	0
51	H	i-pr	3	5-SO ₂ Me	H	H	H	0
59	H	i-pr	3	H	H	H	H	1
60	H		3	H	H	H	H	1
61	H	i-pr	3	5-COOEt	H	H	H	1
62	H	i-pr	3	5-CMe ₂ OH	H	H	H	1
63	H	i-pr	3	6-CH ₂ CHMe ₂	H	H	H	1
64	H	i-pr	3	6-Me	H	H	H	1
65	H	c-pr	3	H	H	H	H	1
66	H	c-pr	3	6-CMe ₂ OH	H	H	H	1
67	H	c-pr	4	H	H	H	H	1
68	H	c-pr	3	5-Br	H	H	H	1
73	H	i-butyl	3	6-CMe ₂ OH	H	H	H	1
74	H	c-pr	3	6-Me	H	H	H	1
75	H	c-pr	3	6-SO ₂ Me	H	H	H	1
76	H	c-pr	3	6-CMe ₂ OH	H	H	5-Br	1
77	H	c-pr	3	6-CMe(CH ₂ OH)OH	H	H	H	1

Table 3



5



Example	R	R ¹	Ar	R ⁴	R ⁶
8	H	i-pr	indol-5-yl	H	H
12	H	i-pr	quinolin-3-yl	H	H
13	H	i-pr	pyrimidin-5-yl	H	H
19	H	c-pr	3-thienyl	H	H
29	H	c-pr	1-oxidopyrimidin-5-yl	H	H
57	H	c-pr		H	H
72	H	i-pr	1-oxidoquinolin-3-yl	H	H

Table 4



10














Example	R	R ¹	Ar	Ar ¹	R ⁴	R ⁶
31	H	i-pr	Ph	4-(pyridin-3-yl)	H	H
58	H	c-pr	Pyridin-3-yl		H	H
69	H	c-pr	Pyridin-3-yl		H	H
70	H	c-pr	1-oxidopyridin-3-yl		H	H
71	H	c-pr	1-oxidopyridin-3-yl		H	H













Table 5



5


R ¹	R ⁴	R ⁶	Ar
c-pr	H	H	

R ¹	R ⁴	R ⁶	Ar
c-pr	H	H	
i-pr	7-Me	H	
i-pr	H	5-Me	
c-pr	H	H	
c-pr	H	H	
c-pr	H	H	
c-Bu	H	6-F	

R^1	R^4	R^6	Ar
	6-F	H	
	H	H	
	5-F	4-F	
i-pr	H	H	
i-pr	H	H	
i-pr	H	H	
c-pr	H	H	
	H	H	

R^1	R^4	R^6	Ar
c-pr	H	H	
c-Bu	H	H	
	H	H	
	H	H	

EXAMPLE 1

5 N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Step 1: Ethyl 3-(3-bromoanilino)-2-(2-chloronicotinoyl) acrylate.

A mixture of ethyl 2-chloronicotinoyl acetate (41.1g, 180.5mmol), triethyl orthoformate (40.12g, 271mmol) and acetic anhydride (92.05g, 902.5mmol) was heated at 130°C for 2.5 hours. The volatile components were distilled off and the residue was co-evaporated twice with xylene. The oily residue was dissolved in methylene chloride (250mL) and 3-bromoaniline (37.25g, 216.6mmol) was added slowly. The resulting solution was stirred at room temperature for 18 hours, and the solvent evaporated away. The resulting crude compound was used as such in the next step.

Step 2: Ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate.

The crude compound from Step 1 was dissolved in tetrahydrofuran (500mL), the solution was cooled to 0°C, and sodium hydride (as a 60% dispersion in oil, 9.4g, 235mmol) was added in portions. After stirring at 0° for 1 hour, the mixture was allowed to warm up to room temperature. After 2 hours, water (400mL) was added to the suspension and the insoluble solid was filtered and washed copiously with water. When dry, the solid was stirred in ether (150mL) at room temperature for 24 hours and filtered to afford the title compound as a cream-colored solid.

¹H NMR (Acetone-d₆) δ 1.32 (t, 3H), 4.29 (q, 2H), 7.54-7.63 (m, 2H), 7.69 (dd, 1H), 7.78 (dd, 1H), 7.93 (s, 1H), 8.66-8.71 (m, 3H).

Step 3: 1-(3-Bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

A suspension of ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from Step 2 (52.5g, 140.7mmol) in a mixture of tetrahydrofuran (400mL), methanol (400mL) and 1N aqueous sodium hydroxide (280mL) was heated at ca 50°C with stirring for 20 minutes. After cooling, the mixture was diluted with water (300mL) and 1N aqueous HCl (325mL) was added. After stirring for 45 minutes, the precipitate was filtered, washed well with water and dried to afford the title acid as a cream-colored solid.

¹H NMR (Acetone-d₆) δ 7.65 (t, 1H), 7.76 (m, 2H), 7.84 (d, 1H), 7.99 (s, 1H), 8.87 (m, 2H), 9.01 (s, 1H).

Step 4: N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

To a suspension of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 3 (26.3g, 76mmol) and triethylamine (23.2g, 230mmol) in

tetrahydrofuran (1000mL) at 0°C was added isobutyl chloroformate (18.85g, 138mmol). After stirring at 0°C for 2 hours, isopropylamine (23g, 390mmol) was added and the mixture was allowed to warm up to room temperature and stirred overnight. The mixture was then partitioned between ethyl acetate and water, the organic phase was dried and evaporated to a solid which was stirred in ether at room temperature for 3 hours and filtered to afford the N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide as a white solid.

¹H NMR (Acetone-d₆) δ 1.25 (d, 6H), 4.17 (m, 1H), 7.59-7.63 (m, 2H), 7.70 (d, 1H), 7.80 (d, 1H), 7.94 (s, 1H), 8.73 (m, 1H), 8.78 (d, 1H), 8.85 (s, 1H), 9.61 (br, NH).

Step 5: N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


A mixture of N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 4, 3-acetylphenylboronic acid (1.2eq.), *trans*-dibromobis(triphenylphosphine)palladium (II) (0.05eq.), toluene (6mL/mmol), ethanol (2mL/mmol) and 2M aqueous sodium carbonate (8eq.) was refluxed for 1 hour under a nitrogen atmosphere. The mixture was diluted with ethyl acetate and the organic phase was washed with water and brine, dried and evaporated. The crude product was chromatographed on silica gel eluting with a gradient of 20-40% ether in methylene chloride to afford the N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide product as a solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 2.65 (s, 3H), 4.28 (m, 1H), 7.47 (m, 2H), 7.55 (t, 1H), 7.65 (m, 2H), 7.80 (m, 2H), 7.95 (dd, 1H), 8.19 (brs, 1H), 8.70 (dd, 1H), 8.81 (dd, 1H), 9.05 (s, 1H), 9.65 (br, NH).

EXAMPLE 2

N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Step 1: Anion of 4-amino-3,5-dichloropyridine.

10 A suspension of sodium hydride as 60% dispersion in oil (360mg, 9mmol) in tetrahydrofuran (15mL) was cooled to 0°C. A solution of 4-amino-3,5-dichloropyridine (978mg, 6mmol) in tetrahydrofuran (15mL) was added slowly. The resulting mixture was kept at 0° for 2.5 hours.

15 Step 2: Acid chloride of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid.

A suspension of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 3 of Example 1 (690mg, 2mmol) in tetrahydrofuran (12mL) was cooled to 0°C, and oxalyl chloride (381mg, 3mmol) was added, followed by 2 drops of N,N-dimethylformamide. The resulting mixture was then stirred at room temperature for 1 hour then
20 refluxed for 45 minutes and cooled to room temperature.

Step 3: N-(2,6-Dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

The mixture from Step 2 above, as a brown suspension, was added via syringe to the cold suspension of Step 1. The resulting mixture was stirred at room temperature for 18 hours, quenched with aqueous saturated ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from evaporation of the organic phase was triturated with ether (50mL) and filtered, affording the N-(2,6-Dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide as a beige solid.

¹H NMR (Acetone-d₆) δ 7.61-7.70 (m, 2H), 7.76 (d, 1H), 7.81 (d, 1H), 8.00 (s, 1H), 8.62 (s, 2H), 8.80 (br s, 1H), 8.86 (d, 1H), 8.99 (s, 1H), 12.1 (br, NH).


Step 4: N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

Following the procedure of Step 5 of Example 1, but substituting N-(2,6-dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 3 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 2.65 (s, 3H), 7.47 (d, 1H), 7.50-7.60 (m, 2H), 7.70 (m, 2H), 7.82 (d, 2H), 7.98 (d, 1H), 8.20 (s, 1H), 8.55 (s, 2H), 8.75 (brs, 1H), 8.92 (dd, 1H), 9.14 (s, 1H), 12.08 (br, NH).

EXAMPLE 3

N-Isopropyl-1-[3-(4-n-propylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




5

Following the procedure of Step 5 of Example 1, but substituting 4-n-propylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a white solid.

10 ^1H NMR (Acetone- d_6) δ 0.93 (t, 3H), 1.24 (d, 6H), 1.65 (m, 2H), 2.62 (t, 2H), 4.18 (m, 1H), 7.31 (d, 2H), 7.58-7.61 (m, 2H), 7.68-7.72 (m, 3H), 7.87 (d, 1H), 7.95 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.66 (br, NH).

EXAMPLE 4

N-Isopropyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




5

Following the procedure of Step 5 of Example 1, but substituting 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

10 ^1H NMR (Acetone- d_6) δ 1.25 (d, 6H), 2.61 (s, 3H), 4.17 (m, 1H), 7.59 (m, 1H), 7.70 (d, 1H), 7.76 (t, 1H), 7.92 (d, 2H), 7.97 (d, 1H), 8.07-8.10 (m, 3H), 8.72 (brs, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.65 (br, NH).

EXAMPLE 5

N-Isopropyl-1-[3-(2-methylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



5


Following the procedure of Step 5 of Example 1, but substituting 2-methylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

10 ^1H NMR (Acetone- d_6) δ 1.24 (d, 6H), 2.35 (s, 3H), 4.17 (m, 1H), 7.27-7.34 (m, 4H), 7.56-7.60 (m, 2H), 7.65 (m, 2H), 7.70 (t, 1H), 8.74 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.64 (br, NH).

EXAMPLE 6

N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5

Step 1: N-Isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

10 Following the procedure of Example 1, Step 4, but substituting N-isopropyl-N-methylamine for isopropylamine the N-Isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained in as a yellow solid.

¹H NMR (Acetone-d₆) (Appears as two rotamers of the amide) δ 1.18 (m, 6H), 2.85 (s, 3H), 4.05 (m, 0.5H), 4.84 (m, 0.5H), 7.49-7.64 (m, 3H), 7.72 (d, 1H), 7.86 (s, 1H), 8.14 (s, 1H), 8.65 (d, 2H).

15

Step 2: N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


20 Following the procedure of Step 5 of Example 1, but substituting N-isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

¹H NMR (CDCl₃) (Appears as two rotomers of the amide) δ 1.23 (m, 6H), 2.62 (s, 3H), 4.00 (m, 0.5H), 4.92 (m, 0.5H), 7.38-7.55 (m, 2H), 7.63-7.77 (m, 5H), 8.03 (d, 2H), 8.14 (s, 0.5H), 8.21 (s, 0.5H), 8.65 (m, 1H), 8.75-8.80 (m, 1H).

5

EXAMPLE 7

N-Isopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



10


Following the procedure of Step 5 of Example 1, but substituting pyridine-3-boronic acid 1,3-propanediol cyclic ester for 3-acetylphenylboronic acid and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) the title compound was obtained as a beige solid.

15

¹H NMR (Acetone-d₆) δ 1.24 (d, 6H), 4.17 (m, 1H), 7.48 (m, 1H), 7.60 (m, 1H), 7.71 (dd, 1H), 7.78 (t, 1H), 7.95 (dd, 1H), 8.05 (brs, 1H), 8.15 (m, 1H), 8.60 (m, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 8.99 (brs, 1H), 9.65 (br, NH).

EXAMPLE 8

N-Isopropyl-1-[3-(indol-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




5

Following the procedure of Step 5 of Example 1, but substituting 5-indolylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as an off-white solid.

10 ^1H NMR ((DMSO- d_6) δ 1.20 (d, 6H), 4.10 (m, 1H), 6.47 (s, 1H), 7.38 (brs, 1H), 7.46-7.52 (m, 3H), 7.59-7.66 (m, 2H), 7.87-7.93 (m, 3H), 8.72-8.81 (m, 3H), 9.67 (br, NH), 11.2 (br, NH).

EXAMPLE 9

N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5

Step 1: N-tert-Butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 1, Step 4, but substituting tert-butylamine for isopropylamine the N-tert-Butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained as a yellow solid.

¹H NMR (Acetone-d₆) δ 1.44 (s, 9H), 7.58-7.62 (m, 2H), 7.70 (dd, 1H), 7.78 (dd, 1H), 7.93 (br s, 1H), 8.72 (m, 1H), 8.77 (dd, 1H), 8.81 (s, 1H), 9.73 (br, NH).


15 Step 2: N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 5 of Example 1, but substituting N-tert butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained in 93% yield as a white solid.

¹H NMR (Acetone-d₆) δ 1.45 (s, 9H), 2.61 (s, 3H), 7.59 (m, 1H), 7.69-7.72 (m, 1H), 7.77 (t, 1H), 7.92-7.99 (m, 3H), 8.07-8.11 (m, 3H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.91 (s, 1H), 9.79 (br, NH).

EXAMPLE 10

5 N-(2,6-Dichloropyridin-4-yl)-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide




10 Following the procedure of Step 4 of Example 2 but substituting [1,1'-bis
(diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-
dibromobis(triphenylphosphine)palladium (II) and pyridine-3-boronic acid 1,3-propanediol
cyclic ester for 3-acetylphenylboronic acid the title compound was obtained as a glassy solid.

¹H NMR (Acetone-d₆) δ 7.48 (m, 1H), 7.68 (m, 1H), 7.77-7.82 (m, 2H), 7.98 (m,
1H), 8.12-8.17 (m, 2H), 8.61 (m, 1H), 8.62 (s, 2H), 8.80 (m, 1H), 8.88 (dd, 1H), 8.99 (brs, 1H),
15 9.06 (s, 1H), 12.2 (br, NH).

EXAMPLE 11

N-Isopropyl-1-{3-[4-(4-tertbutyloxycarbonylpiperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Step 1: 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine

10 To a suspension of 1-(4-bromophenyl)piperazine hydrochloride (103.15g, 371.59mmol) in acetonitrile (1.5 L) at 0°C under a nitrogen atmosphere was added a catalytic amount of 4-dimethylaminopyridine (4.54g, 37.159mmol) followed by triethylamine (155mL, 1114.77mmol) and di-*tert*-butyl dicarbonate (121.65g, 557.385mmol, dissolved in a minimum amount of acetonitrile) and the resulting reaction mixture was warmed to room temperature and stirred for 5.5 hours. The reaction mixture was filtered, ethyl acetate was added and the organic phase was washed with 10 % aqueous citric acid, water (2X) and brine, then dried and

15 evaporated to afford the crude 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine product which was used as such in the next step.

Step 2: 3-(4-tert-Butyloxycarbonylpiperazin-1-yl)phenylboronic acid

20 To the 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine from Step 1 (118.30g, 346.9mmol) in tetrahydrofuran /toluene (1/1, 1.5 L) at -78°C under nitrogen was

added n-butyllithium (2.5M, 160mL, 398.9mmol) dropwise and the resulting reaction mixture was stirred at -78°C for 20 minutes. Triisopropyl borate (96.1mL, 416.3mmol) was added dropwise and the reaction was warmed to 0°C and stirred for 2 hours. Aqueous saturated ammonium chloride (400mL), water (100mL) and 1 equivalent of H_3PO_4 (20mL) were added and the mixture was stirred for 15 minutes and then concentrated to a volume of approximately 200mL (at which stage the mixture became bluish and a precipitate formed). The mixture was slowly diluted with heptane (800mL) and the resulting suspension was stirred overnight. The suspension was filtered, the solid was washed with heptane and dried to afford the title boronic acid.


Step 3: N-Isopropyl-1-{3-[4-(4-tertbutyloxycarbonylpiperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 5 of Example 1 but substituting [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) and the boronic acid from Step 2 above for 3-acetylphenylboronic acid the 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine compound was obtained as a solid.

^1H NMR (CDCl_3) δ 1.30 (d, 6H), 1.49 (s, 9H), 3.18 (m, 4H), 3.58 (m, 4H), 4.29 (m, 1H), 6.98 (d, 2H), 7.32 (d, 1H), 7.45 (m, 1H), 7.53 (d, 2H), 7.55-7.62 (m, 2H), 7.72 (d, 1H), 8.70 (m, 1H), 8.82 (d, 1H), 9.07 (s, 1H), 9.68 (br, NH).

EXAMPLE 12

N-Isopropyl-1-[3-(quinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide




Following the procedure of Step 5 of Example 1, but substituting 3-quinolineboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 4.29 (m, 1H), 7.49 (m, 2H), 7.61 (t, 1H), 7.70-7.78 (m, 3H), 7.86-7.92 (m, 2H), 8.14 (d, 1H), 8.36 (s, 1H), 8.71 (m, 1H), 8.84 (dd, 1H), 9.10 (s, 1H), 9.19 (s, 1H), 9.67 (br, NH).

EXAMPLE 13


10 N-Isopropyl-1-[3-(pyrimidin-5-yl) phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide



15 Following the procedure of Step 5 of Example 1, but substituting 5-pyrimidineboronic acid for 3-acetylboronic acid the title compound was obtained as a solid.

¹H NMR (CDCl₃) δ 1.28 (d, 6H), 4.27 (m, 1H), 7.48 (dd, 1H), 7.52 (m, 1H), 7.65 (s, 1H), 7.74 (m, 2H), 8.68 (m, 1H), 8.72 (d, 1H), 8.98 (s, 2H) 9.03 (s, 1H), 9.22 (s, 1H), 9.62 (br, NH).

EXAMPLE 14

N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamideStep 1: N-Cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 1, Step 4, but substituting cyclopropylamine for isopropylamine the N-Cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained as a fluffy white solid.

^1H NMR (Acetone- d_6) δ 0.59 (m, 2H), 0.80 (m, 2H), 2.96 (m, 1H), 7.59-7.68 (m, 2H), 7.72 (dd, 1H), 7.82 (dd, 1H), 7.97 (s, 1H), 8.72-8.81 (m, 2H), 8.89 (s, 1H), 9.70 (br, NH).


Step 2: N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 7 but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a cream-coloured solid.

^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 7.52 (m, 1H), 7.63-7.69 (m, 2H), 7.74 (t, 1H), 7.97 (d, 1H), 8.07 (brs, 1H), 8.17 (d, 1H), 8.61 (m, 1H), 8.73 (dd, 1H), 8.79 (m, 1H), 8.85 (s, 1H), 8.99 (brs, 1H), 9.74 (br, NH).

EXAMPLE 15

N-Isopropyl-1-[3-(5-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide




Following the procedure of Step 5 of Example 1 but substituting 5-methylthiopyridine-3-boronic acid for 3-acetylphenylboronic acid and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) the title compound was obtained as a solid.

^1H NMR (CDCl_3) δ 1.33 (d, 6H) 2.60 (s, 3H), 4.33 (m, 1H), 7.48-7.54 (m, 2H), 7.66 (m, 1H), 7.73 (t, 1H), 7.78-7.81 (m, 2H), 8.55 (s, 1H), 8.66 (s, 1H), 8.74 (m, 1H), 8.87 (d, 1H), 9.09 (s, 1H), 9.69 (br, NH).

EXAMPLE 16

N-Cyclopropyl-1-[3-(4-hydroxymethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




Following the procedure of Step 2 of Example 14 but substituting 4-hydroxymethylphenyl boronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a solid.

10

^1H NMR (CDCl_3) δ 0.71 (m, 2H), 0.89 (m, 2H), 1.88 (t, 1H), 3.03 (m, 1H), 4.78 (d, 2H), 7.43 (d, 1H), 7.46-7.52 (m, 3H), 7.61-7.69 (m, 4H), 7.80 (d, 1H), 8.73 (m, 1H), 8.83 (dd, 1H), 9.10 (s, 1H), 9.82 (br, NH).

EXAMPLE 17

N-Cyclopropyl-1-[3-(pyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



5

Following the procedure of Step 5 of Example 1 but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-pyridineboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a white solid.


10

^1H NMR ((DMSO- d_6) δ 0.57 (m, 2H), 0.77 (m, 2H), 2.90 (m, 1H), 7.64 (m, 1H), 7.72-7.89 (m, 4H), 8.03 (d, 1H), 8.13 (s, 1H), 8.66-8.78 (m, 4H), 8.84 (s, 1H), 9.72 (br, NH).

EXAMPLE 18

N-Cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




Following the procedure of Step 2 of Example 14 but substituting 4-ethylthiobenzeneboronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a solid.

10

^1H NMR (CDCl_3) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.48 (t, 3H), 3.03 (m, 3H), 7.42 (d, 3H), 7.50 (m, 1H), 7.57 (d, 2H), 7.64 (s, 1H), 7.68 (t, 1H), 7.78 (d, 1H), 8.75 (m, 1H); 8.85 (d, 1H), 9.10 (s, 1H), 9.83 (br, NH).

EXAMPLE 19

N-Cyclopropyl-1-[3-(3-thienyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




Following the procedure of Step 2 of Example 14 but substituting 3-thiopheneboronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a white solid.

10 ^1H NMR (Acetone- d_6) δ 0.60 (m, 2H), 0.79 (m, 2H), 2.96 (m, 1H), 7.57-7.72 (m, 5H), 7.92-7.98 (m, 2H), 8.05 (s, 1H), 8.74 (s, 1H), 8.78 (d, 1H), 8.93 (s, 1H), 9.74 (br, NH).

EXAMPLE 20

N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5

Step 1: 4-Sulfamoylbenzeneboronic acid pinacol ester

10 A mixture of 4-bromobenzenesulfonamide, diboron pinacol ester (1.1eq), potassium acetate (3.5eq) and 1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-dimethylformamide (4ml/mmol) was heated at 85°C for 18 hours. After quenching with saturated aqueous ammonium chloride solution the mixture was partitioned between ethyl acetate and water and the product from the organic phase was chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and hexane to afford the 4-

15 Sulfamoylbenzeneboronic acid pinacol ester as a solid.

Step 2: N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


20 A mixture of N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, boronate from Step 1 (1.2eq), palladium acetate (0.1eq), triphenylphosphine (0.35eq) and 2M aqueous sodium carbonate (3.5eq) in n-propanol (10ml/mmol) was stirred at 85°C for 1 hour. After cooling, the mixture was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water, and the product from the organic phase was chromatographed on silica gel eluting with a 1:5:4 mixture of ethanol, ethyl

acetate and methylene chloride to afford the N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

¹H NMR (Acetone-d₆) δ 0.62 (m, 2H), 0.82 (m, 2H), 2.98 (m, 1H), 6.66 (br, NH₂), 7.64 (m, 1H), 7.74 (m, 1H), 7.80 (t, 1H), 7.97-8.05 (m, 5H), 8.10 (m, 1H), 8.76 (m, 1H), 8.81 (dd, 1H), 8.97 (s, 1H), 9.77 (br, NH).

EXAMPLE 21

N-Isopropyl-1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: Ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure of Step 5 of Example 1, but substituting ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from Step 2 of Example 1 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, and 3-ethoxybenzeneboronic acid for 3-acetylbenzeneboronic acid, the Ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate compound was obtained as a solid.

Step 2: 1-[3-(3-Ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

Following the procedure of Step 3 of Example 1 but substituting ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from step 1 for ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate the 1-[3-(3-Ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained and used without purification in the next step.


Step 3: N-Isopropyl- 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5 A mixture of 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 2 and thionyl chloride (4eq) in tetrahydrofuran (10ml/mmol) was refluxed for 45 minutes, then evaporated. The residue was dissolved in the same volume of tetrahydrofuran, isopropylamine (5eq) was added and the mixture was stirred at room temperature for 18 hours. After quenching with saturated aqueous ammonium chloride solution, the resulting mixture was partitioned between ethyl acetate and water, and the product from the organic phase was chromatographed on silica gel eluting with 10% ether in methylene chloride to afford the N-Isopropyl- 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 1.42 (t, 3H), 4.08 (q, 2H), 4.28 (m, 1H), 6.91 (d, 1H), 7.12 (s, 1H), 7.18 (d, 1H), 7.34 (t, 1H), 7.40 (d, 1H), 7.46 (m, 1H), 7.60-7.65 (m, 2H), 7.75 (d, 1H), 8.71 (brs, 1H), 8.82 (dd, 1H), 9.08 (s, 1H), 9.70 (br, NH).

EXAMPLE 22

N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: Ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure of Step 5 of Example 1, but substituting ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from step 2 of example 1 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, and 4-methylthiobenzenboronic acid for 3-acetylbenzenboronic acid, the Ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate compound was obtained as a solid.

Step 2: 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

Following the procedure of Step 3 of Example 1 but substituting ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from Step 1 for ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate the 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a solid.


Step 3: N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 21 but substituting 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid for 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid the N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

^1H NMR (Acetone- d_6) δ 1.24 (d, 6H), 2.52 (s, 3H), 4.18 (m, 1H), 7.37 (d, 2H), 7.58-7.62 (m, 2H), 7.69-7.73 (m, 3H), 7.87 (d, 1H), 7.96 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.91 (s, 1H), 9.65 (br, NH).

EXAMPLE 23

N-Isopropyl-1-[3-(3-acetyl-4-hydroxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide




A mixture of 5'-bromo-2'-hydroxyacetophenone, diboron pinacol ester (1.25eq), potassium acetate (3eq) and [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-dimethylformamide (10ml/mmol) was stirred at 80°C for 3 hours and cooled down. A solution of N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 1, Step 4 (0.75eq) in N,N-dimethylformamide (7ml/mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) and 2M aqueous sodium carbonate (8.5eq) were added and the resulting mixture was stirred at 80°C for 2.5 hours. The cooled mixture was partitioned between ethyl acetate and water and the product from the organic phase was chromatographed on silica gel eluting with 60% ethyl acetate in hexane to afford the title compound as a light yellow solid.

¹H NMR (Acetone-d₆) δ 1.24 (d, 6H), 2.75 (s, 3H), 4.19 (m, 1H), 7.06 (d, 1H), 7.59-7.63 (m, 2H), 7.72 (t, 1H), 7.92 (d, 1H), 7.97 (d, 1H), 8.02 (s, 1H), 8.33 (s, 1H), 8.73 (m, 1H), 8.78 (dd, 1H), 8.90 (s, 1H), 9.65 (br, NH).

EXAMPLE 24

N-Isopropyl-1-[3-(5-carboethoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




Following the procedure of Example 23 but substituting N-isopropyl-1-(3-

10 bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for 5'-bromo-2'-
hydroxyacetophenone and ethyl 5-bromonicotinate for N-isopropyl-1-(3-bromophenyl)-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide the title compound was obtained as a beige
solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 1.40 (t, 3H), 4.28 (m, 1H), 4.42 (q, 2H), 7.45-
7.51 (m, 2H), 7.68 (s, 1H), 7.71 (t, 1H), 7.80 (d, 1H), 8.49 (s, 1H), 8.59 (m, 1H), 8.82 (d, 1H),
15 9.03 (s, 1H), 9.07 (s, 1H), 9.23 (s, 1H), 9.64 (br, NH).

EXAMPLE 25

N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 3-Bromo-5-(1-hydroxy-1-methylethyl)pyridine


To a solution of ethyl 5-bromonicotinate (1.02g, 4.4mmol) in diethyl ether (15ml) at -30°C was added a 3M solution of methyl magnesium bromide (4ml, 12mmol) in ether. The resulting slurry was then refluxed for 2 hours then cooled and quenched with an excess of 0.5M aqueous monobasic sodium phosphate and partitioned between ether and water. The product from the organic phase was chromatographed on silica gel eluting with a 2:1:2 mixture of ether, pentane and ammonia-saturated methylene chloride to afford the 3-Bromo-5-(1-hydroxy-1-methylethyl)pyridine compound as a yellow oil.

Step 2: N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

Following the procedure of Example 24, but substituting the 3-bromo-5-(1-hydroxy-1-methylethyl)pyridine from Step 1 for ethyl 5-bromonicotinate, the title compound was obtained as a yellow foam.

^1H NMR (CDCl_3) δ 1.28 (d, 6H), 1.62 (s, 6H), 2.52 (brs, 1H), 4.25 (m, 1H), 7.41-7.48 (m, 2H), 7.60-7.68 (m, 2H), 7.75 (d, 1H), 8.05 (s, 1H), 8.67-8.71 (m, 3H), 8.80 (dd, 1H), 9.03 (s, 1H), 9.66 (br, NH).

EXAMPLE 26

N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamideStep 1: 5-Bromo-2-(2-methylpropyl)pyridine

To a solution of 2,5-dibromopyridine (4.5g, 19mmol) in tetrahydrofuran (50ml) was added [1,1'-bis (diphenylphosphino)ferrocene]dichloronickel (II) (103mg, 0.19mmol) and the resulting mixture was cooled to -10°C . A 2M solution of isobutylmagnesium bromide in ether (12.4ml, 24.7mmol) was added slowly and the mixture was stirred at -10 to 10°C for 3.5 hours. After quenching with saturated aqueous ammonium chloride solution, the mixture was partitioned between ether and water and the product from the organic phase was chromatographed on silica gel eluting with 10% ether in pentane to afford the 5-Bromo-2-(2-methylpropyl)pyridine compound as a volatile oil.


Step 2: N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 24 but substituting 5-bromo-2-(2-methylpropyl)pyridine from Step 1 for ethyl 5-bromonicotinate the N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

^1H NMR (CDCl_3) δ 0.92 (d, 6H), 1.28 (d, 6H), 2.10 (m, 1H), 2.69 (d, 2H), 4.28 (m, 1H), 7.19 (d, 1H), 7.40-7.47 (m, 2H), 7.60 (s, 1H), 7.64 (t, 1H), 7.73 (d, 1H), 7.79 (dd, 1H), 8.68 (m, 1H), 8.77-8.83 (m, 2H), 9.05 (s, 1H), 9.66 (br, NH).

EXAMPLE 27

5 N-Isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide



10 Step 1: 3-Acetyl-5-bromopyridine

To a solution of ethyl 5-bromonicotinate (3.9g, 16.9mmol) in ether (50ml) at 0°C was added a 3M solution of methylmagnesium bromide (16.9ml, 50.8mmol). The resulting thick slurry was warmed slowly to room temperature and after 1.5 hours it was poured slowly into an excess of 1M aqueous monobasic sodium phosphate. The mixture was partitioned between ether and water and the product from the organic phase was chromatographed on silica gel, eluting with a 1:1:2 mixture of ether, pentane and ammonia-saturated methylene chloride to afford the 3-acetyl-5-bromopyridine compound. This preparation also afforded 3-bromo-5-(1-hydroxy-1-methylethyl)pyridine described in Example 25.


20 Step 2: N-Isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

Following the procedure of Example 23 but substituting 3-acetyl-5-bromopyridine from Step 1 for ethyl 5-bromonicotinate the N-isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 2.69 (s, 3H), 4.28 (m, 1H), 7.48 (dd, 1H), 7.51 (d, 1H), 7.69 (s, 1H), 7.72 (t, 1H), 7.80 (d, 1H), 8.42 (s, 1H), 8.69 (m, 1H), 8.82 (d, 1H), 9.05 (s, 2H) 9.17 (s, 1H), 9.63 (br, NH).

EXAMPLE 28

N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Step 1: 5-Bromo-2-methylpyridine

Following the procedure of Step 1 of Example 26 but substituting methylmagnesium chloride for isobutylmagnesium bromide the 5-bromo-2-methylpyridine compound was obtained as a solid.


Step 2: N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 24 but substituting 5-bromo-2-methylpyridine from Step 1 for ethyl 5-bromonicotinate the N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 1.32 (d, 6H), 2.63 (m, 3H), 4.30 (m, 1H), 7.25 (d, 1H), 7.45-7.51 (m, 2H), 7.63 (s, 1H), 7.69 (t, 1H), 7.77 (d, 1H), 7.82 (dd, 1H), 8.72 (m, 1H), 8.78 (s, 1H), 8.85 (d, 1H), 9.08 (s, 1H), 9.68 (br, NH).

EXAMPLE 29

N-Cyclopropyl-1-[3-(1-oxidopyrimidin-5-yl) phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide



Step 1: 5-Bromo-1-oxidopyrimidine

To 5-bromopyrimidine (2.05g, 12.9mmol) in methylene chloride (25ml) was added m-chloroperoxybenzoic acid (ca 70% pure, 3.17g, 12.9mmol) and the resulting mixture was stirred at room temperature for 5 days. Calcium hydroxide (1g) was added and after 10 minutes the mixture was filtered through celite. The product from evaporation of the filtrate was chromatographed on silica gel eluting with ethyl acetate to afford the 5-bromo-1-oxidopyrimidine compound as a white solid.


Step 2: N-Cyclopropyl-1-[3-(1-oxidopyrimidinyl-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Example 24 but substituting 5-bromo-1-oxidopyrimidine from Step 1 for ethyl 5-bromonicotinate and N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(1-oxidopyrimidinyl-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

^1H NMR (CDCl_3) δ 0.66 (m, 2H), 0.84 (m, 2H), 2.97 (m, 1H), 7.48 (m, 1H), 7.58 (d, 1H), 7.65 (s, 1H), 7.71 (d, 1H), 7.77 (t, 1H), 8.46 (s, 1H), 8.60 (s, 1H), 8.68 (brs, 1H), 8.81 (dd, 1H), 8.98 (s, 1H), 9.02 (s, 1H), 9.72 (br, NH).

EXAMPLE 30

1-{3-[6-(1-Hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine

To a suspension of 2,5-dibromopyridine in toluene (12ml/mmol) cooled to -78°C was added n-butyllithium 2.5M in hexanes (1.05eq) and the resulting mixture was stirred in the cold for 2.5 hours. Acetone (2eq) was added and stirring was continued for 1.5h. After quenching with saturated aqueous ammonium chloride solution, the mixture was warmed to room temperature and partitioned between ethyl acetate and water. The product from the organic phase was chromatographed on silica gel eluting with 20% ethyl acetate in hexane to afford the 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine compound as a syrup.

Step 2: 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide

To a solution of 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine from Step 1 in methylene chloride (5ml/mmol) at room temperature was added m-chloroperoxybenzoic acid 70% (1.1eq) and the resulting mixture was stirred at room temperature for 18 hours. An excess

of calcium hydroxide was added and after 5 minutes the mixture was filtered through a bed of celite. The crude product from evaporation of the filtrate was chromatographed on silica gel eluting with 80% ethyl acetate in hexane and the 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide compound was obtained as a white solid.

5
Step 3: 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 4 of Example 1 but substituting 28% aqueous ammonium hydroxide for isopropylamine the 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

10
Step 4: 1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Example 24 but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from Step 2 above for ethyl 5-bromonicotinate and 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the 1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 1.76 (s, 6H), 5.83 (br, 1H, NH). 7.50 (d, 1H), 7.55 (m, 1H), 7.57-7.62 (m, 2H), 7.65 (m, 2H), 7.72-7.78 (m, 2H), 8.55 (s, 1H, OH), 8.75 (m, 1H), 8.90 (dd, 1H), 9.08 (s, 1H), 9.52 (br, 1H, NH).

EXAMPLE 31

N-Isopropyl-1-[3-[4-(pyridin-3-yl)phenyl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5

Step 1: N-Isopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

10

A mixture of N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 4 of Example 1, diboron pinacol ester (1.1eq), potassium acetate (3.5eq) and [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-dimethylformamide (5ml/mmol) was stirred at 85°C for 18 hours. A further amount of diboron pinacol ester (0.4eq) and palladium catalyst (0.05eq) were added and heating and stirring were continued for a further 24 hours. After cooling, the mixture was partitioned between ethyl acetate and water, and the crude product from the organic phase was chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and hexane. The product was then stirred in hexane at room temperature for several hours and filtered to afford the N-Isopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a white solid.

20

Step 2: 3-(4-Bromophenyl)pyridine

A mixture of pyridine-3-boronic acid 1,3-propanediol cyclic ester, 4-bromoiodobenzene (1.1eq), [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II)

(0.05eq) and 2M aqueous sodium carbonate (5eq) in N,N-dimethylformamide (2ml/mmol) was stirred at 85°C for 4 hours. After quenching with saturated aqueous ammonium chloride solution, the mixture was partitioned between ethyl acetate and water, and the crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ethyl acetate and hexane to afford the 3-(4-Bromophenyl)pyridine compound as a solid.


Step 3: N-Isopropyl-1-{3-[4-(pyridin-3-yl)phenyl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

A mixture of the boronate from Step 1, 3-(4-bromophenyl)pyridine from Step 2 (1.5eq), [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) and 2M aqueous sodium carbonate (5eq) in N,N-dimethylformamide (7ml/mmol) was stirred at 85°C for 1 hour. After cooling, the mixture was partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 7:3 mixture of ethyl acetate and methylene chloride to afford the N-Isopropyl-1-{3-[4-(pyridin-3-yl)phenyl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

¹H NMR (CDCl₃) δ 1.30 (d, 6H), 4.25 (m, 1H), 7.35 (m, 1H), 7.39-7.48 (m, 2H), 7.60-7.75 (m, 6H), 7.80 (d, 1H), 7.90 (d, 1H), 8.58 (d, 1H), 8.70 (m, 1H), 8.82 (d, 1H), 8.88 (s, 1H), 9.08 (s, 1H), 9.68 (br, NH).

EXAMPLE 32

N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: N-Cyclopropyl-1-[3-(4,4,5,5,-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 1 of Example 31 but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 of example 14 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide the N-Cyclopropyl-1-[3-(4,4,5,5,-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

Step 2: 3-Bromo-5-methylsulfonylpyridine

To 3,5-dibromopyridine (2.96g, 12.5mmol) in diethyl ether (70ml) at -78°C was added n-butyllithium 1.6M in hexanes (8.6ml, 13.7mmol) and the resulting mixture was stirred in the cold for 3 hours. Dimethyl disulfide (1.12ml, 12.5mmol) was added and the mixture was warmed to room temperature, then partitioned between ether and water. To the crude product from evaporation of the organic phase was added tetrahydrofuran (80ml), methanol (20ml), oxone (17g) and enough saturated aqueous sodium bicarbonate to afford a slightly basic medium. After stirring for 4 hours at room temperature, an excess of 1M aqueous sodium metabisulfite was added, the organic solvents were evaporated, and the residue was partitioned between ethyl acetate and water. The crude product from the organic phase was stirred in a small volume of ethyl acetate and filtered to afford the 3-Bromo-5-methylsulfonylpyridine compound as a solid.


Step 3: N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 31 but substituting 3-bromo-5-methylsulfonylpyridine from Step 2 above for 3-(4-bromophenyl)pyridine, and N-cyclopropyl-1-[3-(4,4,5,5,-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 1 for N-isopropyl-1-[3-(4,4,5,5,-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 0.71 (m, 2H), 0.90 (m, 2H), 3.03 (m, 1H), 3.21 (s, 3H), 7.53 (m, 1H), 7.60 (d, 1H), 7.74 (s, 1H), 7.80 (t, 1H), 7.86 (d, 1H), 8.45 (m, 1H), 8.74 (m, 1H), 8.86 (d, 1H), 9.09 (s, 1H), 9.20 (d, 2H), 9.78 (br, NH).

EXAMPLE 33

N-Cyclopropyl-1-{3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: Methyl 2-bromoisonicotinate

To a solution of 2-bromoisonicotinic acid (*Chem. Pharm. Bull.*, **38**:2446(1990)) (2.0g) in tetrahydrofuran (100ml) was added excess ethereal diazomethane and the resulting mixture was stirred at room temperature for 1 hour. The mixture was evaporated and the product chromatographed on silica gel eluting with a 1:3 mixture of ethyl acetate and hexane to afford the Methyl 2-bromoisonicotinate ester as a colorless liquid.

Step 2: 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine

Following the procedure of Step 1 of Example 25, but substituting methyl 2-bromoisonicotinate from Step 1 for ethyl 5-bromonicotinate, the 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine compound was obtained as a white solid.

Step 3: 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide

Following the procedure of Step 2 of Example 30 but substituting 2-bromo-4-(1-hydroxy-1-methylethyl)pyridine from Step 2 for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine the 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide compound was obtained as a white solid.


Step 4: N-Cyclopropyl-1-{3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 2-bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide from Step 3 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

¹H NMR (DMSO-d₆) δ 0.57 (m, 2H), 0.79 (m, 2H), 1.45 (s, 6H), 2.90 (m, 1H), 5.35 (s, 1H, OH), 7.48 (m, 1H), 7.64 (m, 1H), 7.72 (m, 3H), 8.11 (m, 2H), 8.30 (d, 1H), 8.72 (dd, 1H), 8.78 (m, 1H), 8.82 (s, 1H), 9.72 (br, NH).

EXAMPLE 34

N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine

A solution of 2,5-dibromopyridine in diethyl ether (5ml/mmol) was cooled to -78°C, and n-butyllithium 2.5M in hexanes (1.05eq) was added slowly. After 2h in the cold, acetone (1.3eq) was added and stirring was continued for 1 hour. The resulting mixture was quenched with saturated aqueous ammonium chloride solution, warmed to room temperature, and partitioned between ether and water. The crude product from the organic phase was

trituated with 1:1 ether-hexane and filtered to afford the 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine compound as a solid.


Step 2: N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 0.71 (m, 2H), 0.90 (m, 2H), 1.68 (s, 6H), 1.85 (s, 1H, OH), 3.04 (m, 1H), 7.45-7.52 (m, 2H), 7.71 (t, 1H), 7.79 (d, 1H), 7.95 (dd, 1H), 8.16 (s, 1H), 8.20 (d, 1H), 8.72 (m, 1H), 8.80-8.87 (m, 2H), 9.12 (s, 1H), 9.82 (br, NH).

EXAMPLE 35

N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine

Following the sequence described in Steps 1-2 of Example 33, but substituting 4-bromopicolinic acid (*Aust. J. Chem.* 24:390(1971)) for 2-bromoisonicotinic acid in Step 1, the 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine compound was obtained as a white solid.


Step 2: N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.78 (m, 2H), 1.48 (s, 6H), 2.91 (m, 1H), 5.27 (s, 1H, OH), 7.62-7.66 (m, 2H), 7.72-7.79 (m, 2H), 8.01 (m, 1H), 8.10 (s, 1H), 8.58 (d, 1H), 8.73-8.79 (m, 2H), 8.84 (s, 1H), 9.73 (br, NH).

EXAMPLE 36

Synthesis of N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine N-oxide

Following the procedure of Step 2 of Example 30, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine from Step 1 of Example 35 for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine N-oxide compound was obtained as a white solid.


Step 2: N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine-N-oxide from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.78 (m, 2H), 1.62 (s, 6H), 2.90 (m, 1H), 6.99 (s, 1H, OH), 7.65-7.84 (m, 4H), 7.94 (s, 1H), 8.03 (dd, 1H), 8.15 (s, 1H), 8.38 (d, 1H), 8.73-8.78 (m, 2H), 8.83 (s, 1H), 9.73 (br, NH).

EXAMPLE 37

N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 5-Bromo-2-isopropylthiopyridine

To a mixture of 2,5-dibromopyridine (2.07g, 8.73mmol) and 2-propanethiol (0.97ml, 10.4mmol) in N,N-dimethylformamide (20ml) at 0°C was added portionwise sodium hydride 60% dispersed in oil (450mg, 11.3mmol). The resulting mixture was stirred at room temperature for 1 hour, then partitioned between ether and water. The crude product from the organic phase was chromatographed on silica gel eluting with 10% ethyl acetate in hexane to afford the 5-Bromo-2-isopropylthiopyridine compound as a solid.

Step 2: 5-Bromo-2-isopropylsulfonylpyridine


To a solution of 5-bromo-2-isopropylthiopyridine from Step 1 (2.03g, 8.75mmol) in tetrahydrofuran (50ml) and methanol (25ml) at 0°C was added oxone (15.8g, 25.8mmol) and then saturated aqueous sodium bicarbonate (25ml). The resulting mixture was stirred at room temperature for 6 hours. The mixture was quenched with aqueous sodium bicarbonate and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with 20% ethyl acetate in hexane to afford the 5-Bromo-2-isopropylsulfonylpyridine compound as a white solid.

Step 3: N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-isopropylsulfonylpyridine from Step 2 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 0.70 (m, 2H), 0.89 (m, 2H), 1.39 (d, 6H), 3.00 (m, 1H), 3.82 (m, 1H), 7.51 (m, 1H), 7.60 (d, 1H), 7.72 (s, 1H), 7.80 (t, 1H), 7.83 (d, 1H), 8.15-8.24 (m, 2H), 8.72 (m, 1H), 8.86 (dd, 1H), 9.03 (s, 1H), 9.10 (s, 1H), 9.77 (br, NH).

EXAMPLE 38

N-Cyclopropyl-1-[3-(6-methoxypyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Step 1: 5-Bromo-2-methoxypyridine

To a solution of 2,5-dibromopyridine (6.95g; 29mmol) in N,N-dimethylformamide (5ml) was added methanol (3.56ml) and 1M potassium tert-butoxide (32.3ml) and the resulting mixture was stirred at room temperature for 18 hours. The resulting slurry was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ether and hexane to afford the 5-Bromo-2-methoxypyridine compound as an oil.

Step 2: N-Cyclopropyl-1-[3-(6-methoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-methoxypyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-cyclopropyl-1-[3-(6-methoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

^1H NMR (CDCl_3) δ 0.71 (m, 2H), 0.89 (m, 2H), 3.00 (m, 1H), 4.00 (s, 3H), 6.85 (d, 1H), 7.44 (d, 1H), 7.50 (m, 1H), 7.62 (s, 1H), 7.68 (t, 1H), 7.73 (d, 1H), 7.83 (dd, 1H), 8.44 (s, 1H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.82 (br, NH).

EXAMPLE 39

N-Cyclopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-methylpyridine from Step 1 of Example 28 for 3-bromo-5-methylsulfonylpyridine, the title
10 compound was obtained as a solid.

^1H NMR (CDCl_3) δ 0.72 (m, 2H), 0.90 (m, 2H), 2.65 (s, 3H), 3.03 (m, 1H), 7.28 (d, 1H), 7.45-7.53 (m, 2H), 7.66 (s, 1H), 7.72 (t, 1H), 7.80 (d, 1H), 7.84 (dd, 1H), 8.73 (m, 1H), 8.80 (s, 1H), 8.86 (dd, 1H), 9.11 (s, 1H), 9.82 (br, NH).

EXAMPLE 40

N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Step 1: 5-Bromo-2-(2,2,2-trifluoroethoxy)pyridine

Following the procedure of Step 1 of Example 38, but substituting 2,2,2-trifluoroethanol for methanol, with heating at 70°C for 18 hours, the 5-Bromo-2-(2,2,2-trifluoroethoxy)pyridine compound was obtained as an oil.

10

Step 2: N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

15

Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-(2,2,2-trifluoroethoxy)pyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.


20

¹H NMR (CDCl₃) δ 0.72 (m, 2H), 0.90 (m, 2H), 3.03 (m, 1H), 4.85 (q, 2H), 7.00 (d, 1H), 7.43-7.53 (m, 2H), 7.62 (s, 1H), 7.69-7.78 (m, 2H), 7.92 (dd, 1H), 8.42 (s, 1H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.80 (br, NH).

EXAMPLE 41

N-Cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




10

Following the procedure of Step 3 of Example 32, but substituting 3,5-dibromopyridine for 3-bromo-5-methylsulfonylpyridine, the title compound was obtained as a white solid.

^1H NMR (DMSO- d_6) δ 0.58 (m, 2H), 0.79 (m, 2H), 2.90 (m, 1H), 7.65 (m, 1H), 7.71-7.77 (m, 2H), 8.03 (d, 1H), 8.14 (s, 1H), 8.49 (s, 1H), 8.74 (brs, 1H), 8.79 (brs, 1H), 8.86 (s, 1H), 9.01 (s, 1H), 9.73 (br, NH).

EXAMPLE 42

N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamideStep 1: 2-Benzyloxy-5-bromopyridine

A mixture of 2,5-dibromopyridine, benzyl alcohol (1.3eq), potassium hydroxide pellets (2.4eq) and dibenzo-18-crown-6 (0.05eq) in toluene (4ml/mmol) was refluxed with azeotropic removal of water for 3 hours. After evaporation of the toluene, the resulting mixture was partitioned between chloroform and water. The crude product from the organic phase was recrystallized from ether-hexane to afford the 2-Benzyloxy-5-bromopyridine compound as a solid.


Step 2: N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 2-benzyloxy-5-bromopyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.79 (m, 2H), 2.91 (m, 1H), 5.42 (s, 2H) 7.00 (d, 1H), 7.32-7.48 (m, 5H), 7.61-7.72 (m, 3H), 7.90 (d, 1H), 7.99 (s, 1H), 8.14 (d, 1H), 8.59 (s, 1H), 8.73-8.84 (m, 3H), 9.73 (br, NH).

EXAMPLE 43

N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxipyridin-3-yl]phenyl}-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 5-Bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide

Following the procedure of Steps 1 and 2 of Example 30, but substituting dicyclopropyl ketone for acetone in Step 1, the 5-Bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide compound was obtained as a solid.

Step 2: N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxipyridin-3-yl]phenyl}-
1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxipyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

^1H NMR (CDCl_3) δ 0.52 (m, 4H), 0.70 (m, 4H), 0.76 (m, 2H), 0.89 (m, 2H), 1.35 (m, 2H), 3.02 (m, 1H), 7.52 (m, 1H), 7.58 (m, 1H), 7.62 (dd, 1H), 7.68 (s, 1H), 7.73-7.80 (m, 3H), 8.15 (br, 1H, OH), 8.49 (s, 1H), 8.72 (m, 1H), 8.85 (dd, 1H), 9.09 (s, 1H), 9.78 (br, NH).

EXAMPLE 44

N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Step 1: 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide

Following the procedure of Step 2 of Example 30, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine from Step 1 of Example 34 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide compound was obtained as a white solid.

Step 2: N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Step 3 of Example 32, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 0.69 (m, 2H), 0.88 (m, 2H), 1.63 (s, 6H), 2.20 (s, 1H, OH), 2.98 (m, 1H), 7.38-7.49 (m, 3H), 7.52 (d, 1H), 7.70 (t, 1H), 7.98-8.04 (m, 2H), 8.50 (s, 1H), 8.69 (m, 1H), 8.80 (dd, 1H), 9.08 (s, 1H), 9.75 (br, NH).

EXAMPLE 45

N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide

5




Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine from Step 1 of Example 30 for 3-bromo-5-methylsulfonylpyridine, the title compound was obtained as a solid.

10

^1H NMR (CDCl_3) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.62 (s, 6H), 3.02 (m, 1H), 4.85 (s, 1H, OH), 7.48-7.53 (m, 3H), 7.68 (s, 1H), 7.73 (t, 1H), 7.80 (d, 1H), 7.95 (dd, 1H), 8.72 (m, 1H), 8.81 (s, 1H), 8.86 (dd, 1H), 9.10 (s, 1H), 9.78 (br, NH).

EXAMPLE 46

N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 5-(3-Aminophenyl)- 2-(1-hydroxy-1-methylethyl) pyridine

Following the procedure of Step 5 of Example 1, but substituting 3-aminophenylboronic acid for 3-acetyl phenylboronic acid and 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine from Step 1 of Example 30 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the 5-(3-Aminophenyl)- 2-(1-hydroxy-1-methylethyl) pyridine compound was obtained as a solid.

Step 2: 1-{3-[6-(1-Hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

Following the procedures of Steps 1-3 of Example 1, but substituting 5-(3-aminophenyl)- 2-(1-hydroxy-1-methylethyl)pyridine for 3-bromoaniline from Step 1 in the First Step, the 1-{3-[6-(1-Hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a solid.

Step 3: N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Step 4 of Example 1, but substituting the acid from Step 2 for 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid, and

isobutylamine for isopropylamine, the N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a cream-colored solid.

¹H NMR (Acetone-d₆) δ 0.98 (d, 6H), 1.53 (s, 6H), 1.88 (m, 1H), 3.26 (t, 2H), 4.66 (s, 1H, OH), 7.60 (m, 1H), 7.69 (d, 1H), 7.76-7.79 (m, 2H), 7.95 (d, 1H), 8.05 (s, 1H), 8.16 (dd, 1H), 8.73 (m, 1H), 8.79 (dd, 1H), 8.90 (s, 1H), 8.94 (s, 1H), 9.83 (br, NH).

EXAMPLE 47

10 N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



15 Step 1: 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

Following the procedures of Steps 1-3 of Example 1, but substituting 3,5-dibromoaniline for 3-bromoaniline in Step 1, the 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a beige solid.

20 Step 2: N-Cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 4 of Example 1, but substituting the 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 1 for 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid, and cyclopropylamine for

isopropylamine, the N-Cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

Step 3: 2-(1-Hydroxy-1-methylethyl)-5-tributylstannylpyridine

5 To a suspension of 2,5-dibromopyridine in toluene (5ml/mmol) at -78°C, was added n-butyllithium 2.5M in hexanes (1eq) and the resulting mixture was stirred in the cold for 2.5 hours. Acetone (1eq) was added, and the mixture was warmed to -50°C and became a brown solution. After cooling down to -78°C, more n-butyllithium (1eq) was added along with ether (2ml/mmol). After stirring in the cold for a further hour, tributyltin chloride (1.1eq) was added and the mixture was warmed to room temperature and stirred for 2 hours. The mixture was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ethyl acetate and hexane to afford the 2-(1-Hydroxy-1-methylethyl)-5-tributylstannylpyridine compound as a colorless liquid.


Step 4: N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

A mixture of N-cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 2, 2-(1-hydroxy-1-methylethyl)-5-tributylstannylpyridine from Step 3 (1.4eq), 1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq), and cuprous iodide (0.05eq) in N,N-dimethylformamide (15ml/mmol) was stirred at 85°C for 5 hours. After cooling the resulting mixture was partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:6:3 mixture of ethanol, ethyl acetate and methylene chloride to afford the N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

¹H NMR (CDCl₃) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.62 (s, 6H), 3.02 (m, 1H), 4.76 (s, 1H, OH), 7.50-7.56 (m, 2H), 7.62 (s, 1H), 7.69 (s, 1H), 7.90-7.96 (m, 2H), 8.74 (m, 1H), 8.79 (s, 1H), 8.86 (dd, 1H), 9.07 (s, 1H), 9.74 (br, NH).

EXAMPLE 48

N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 2-(1-Hydroxy-1-methylethyl)-6-tributylstannylpyridine

Following the procedure of Step 3 of Example 47, but substituting 2,6-dibromopyridine for 2,5-dibromopyridine, the 2-(1-Hydroxy-1-methylethyl)-6-tributylstannylpyridine compound was obtained.


Step 2: N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 4 of Example 47, but substituting 2-(1-hydroxy-1-methylethyl)-6-tributylstannyl pyridine from Step 1 for 2-(1-hydroxy-1-methylethyl)-5-tributylstannylpyridine, the N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

$^1\text{H NMR}$ (CDCl_3) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.61 (s, 6H), 3.04 (m, 1H), 5.13 (s, 1H, OH), 7.40 (d, 1H), 7.46-7.53 (m, 2H), 7.70-7.76 (m, 2H), 7.85 (t, 1H), 8.13 (s, 1H), 8.22 (d, 1H), 8.73 (m, 1H), 8.87 (d, 1H), 9.12 (s, 1H), 9.83 (br, NH).

EXAMPLE 49

N-Isopropyl-1-[3-(4-methylsulfonylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




To a mixture of N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 22 in tetrahydrofuran (24ml/mmol), methanol (12ml/mmol), and water (12ml/mmol), was added oxone (2.24eq) and the resulting mixture was stirred at room temperature for 2 hours. The mixture was quenched with saturated aqueous sodium bicarbonate and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with 30% ether in methylene chloride to afford the title compound as a white solid.

¹H NMR (Acetone-d₆) δ 1.25 (d, 6H), 3.16 (s, 3H), 4.18 (m, 1H), 7.60 (m, 1H), 7.74 (d, 1H), 7.79 (t, 1H), 7.99 (d, 1H), 8.05 (s, 4H), 8.09 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.93 (s, 1H), 9.64 (br, NH).

EXAMPLE 50

N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



Step 1: 5-Bromo-2-methylthiopyridine

A mixture of 2,5-dibromopyridine and sodium thiomethoxide (1.3eq) in N,N-dimethylformamide (2ml/mmol) was stirred at room temperature for 20 minutes then cooled to 0°C. After diluting with cold water the precipitate was filtered to afford the 5-Bromo-2-methylthiopyridine compound as a solid.

Step 2: N-Cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-methylthiopyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

Step 3: N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of Example 49, but substituting N-cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 2 for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-

carboxamide, the N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

¹H NMR (CDCl₃) δ 0.66 (m, 2H), 0.84 (m, 2H), 2.97 (m, 1H), 3.26 (s, 3H), 7.48 (m, 1H), 7.55 (d, 1H), 7.67 (s, 1H), 7.74-7.80 (m, 2H), 8.14-8.19 (m, 2H), 8.68 (m, 1H), 8.81 (dd, 1H), 8.96 (s, 1H), 9.05 (s, 1H), 9.73 (br, NH).

EXAMPLE 51

N-Isopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




Following the procedure of Example 49, but substituting N-isopropyl-1-[3-(5-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 15 for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a solid.

¹H NMR (CDCl₃) δ 1.33 (d, 6H), 3.20 (s, 3H), 4.31 (m, 1H), 7.52 (m, 1H), 7.60 (d, 1H), 7.73 (s, 1H), 7.79 (t, 1H), 7.86 (d, 1H), 8.48 (m, 1H), 8.73 (m, 1H), 8.88 (d, 1H), 9.08 (s, 1H), 9.19 (d, 2H), 9.68 (br, NH).

EXAMPLE 52

N-Cyclopropyl-1-[3-(4-ethylsulfonylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide




Following the procedure of Example 49, but substituting N-cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 18 for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a solid.

^1H NMR (CDCl_3) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.35 (t, 3H), 3.02 (m, 1H), 3.18 (q, 2H), 7.48-7.56 (m, 2H), 7.70 (s, 1H), 7.75 (t, 1H), 7.84 (m, 3H), 8.03 (d, 2H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.80 (br, NH).

EXAMPLE 53

N-Cyclopropyl-1-[3-(4-ethylsulfinylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5




To a solution of N-cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 18, in a 1:1 mixture of methylene
chloride and methanol (9ml/mmol), was added at 0°C magnesium monoperoxyphthalate
hexahydrate (MMPP, 0.5molareq) and the resulting mixture was stirred in the cold for 2 hours.
The mixture was quenched with saturated aqueous sodium bicarbonate and partitioned between
methylene chloride and water. The crude product from the organic phase was chromatographed
on silica gel eluting with a 90:9:1 mixture of methylene chloride, ethanol and 28% aqueous
ammonium hydroxide to afford the title compound as a solid.

¹H NMR (CDCl₃) δ 0.68 (m, 2H), 0.85 (m, 2H), 1.15 (m, 3H), 2.80 (m, 1H), 2.94
(m, 1H), 2.98 (m, 1H), 7.45-7.50 (m, 2H), 7.65-7.73 (m, 4H), 7.76-7.82 (m, 3H), 8.71 (m, 1H),
8.83 (dd, 1H), 9.06 (s, 1H), 9.78 (br, NH).

EXAMPLE 54

N-Isopropyl-1-[3-[4-(1-oximidoethyl)phenyl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide




To a solution of N-isopropyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 4 in pyridine (11ml/mmol) at room temperature was added hydroxylamine hydrochloride (2.1eq) and the resulting mixture was stirred for 16 hours. The mixture was filtered through celite and the filtrate evaporated. The residue was dissolved in ethyl acetate, washed with saturated aqueous sodium carbonate and then water, dried and evaporated. The residue was stirred in a small volume of acetone and filtered to afford the title compound as a solid.

^1H NMR (CDCl_3) δ 1.29 (d, 6H), 2.27 (s, 3H), 4.30 (m, 1H), 7.39 (d, 1H), 7.46 (m, 1H), 7.56 (d, 2H), 7.59-7.63 (m, 2H), 7.66 (d, 2H), 7.72 (d, 1H), 8.17 (s, 1H, OH), 8.69 (brs, 1H), 8.82 (d, 1H), 9.10 (s, 1H), 9.71 (br, NH).

EXAMPLE 55


N-Isopropyl-1-{3-[4-(4-piperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



To a solution of N-isopropyl-1-{3-[4-(4-tertbutyloxycarbonylpiperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 11 in methylene chloride (10ml/mmol) was added trifluoroacetic acid (6ml/mmol) and the resulting mixture was stirred at room temperature for 2 hours, then warmed gently for 15 minutes. The mixture was evaporated and the crude product was chromatographed on silica gel eluting with a 9:0.9:0.1 mixture of methylene chloride, methanol and 28% aqueous ammonium hydroxide to afford the title compound as a solid.

¹H NMR (CDCl₃) δ 1.29 (d, 6H), 2.99 (m, 4H), 3.16 (m, 4H), 4.25 (m, 1H), 6.94 (d, 2H), 7.29 (d, 1H), 7.42 (m, 1H), 7.50 (d, 2H), 7.52-7.58 (m, 2H), 7.69 (d, 1H), 8.66 (m, 1H), 8.78 (dd, 1H), 9.04 (s, 1H), 9.69 (br, NH).

EXAMPLE 56

N-Cyclopropyl-1-[3-(4-methylsulfonylmethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamideStep 1: N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

A mixture of N-cyclopropyl-1-[3-(4-hydroxymethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 16, carbon tetrabromide (2eq), and diphos (0.6 molareq) in methylene chloride (15ml/mmol) was stirred at room temperature for 3 hours. The mixture was concentrated at room temperature and chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and methylene chloride to afford the N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound.


Step 2: N-Cyclopropyl-1-[3-(4-methylsulfonylmethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

To a solution of N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 1 in N,N-dimethylformamide (20ml/mmol) was added methanesulfinic acid sodium salt (1.3eq) and the resulting mixture was stirred at room temperature for 18 hours. To the mixture was added saturated aqueous ammonium chloride solution and ethyl acetate, and the insoluble solid was filtered and washed well with water, hexane, ether and ethyl acetate to afford the title compound as a solid.

¹H NMR (CDCl₃) δ 0.72 (m, 2H), 0.89 (m, 2H), 2.85 (s, 3H), 3.04 (m, 1H), 4.34 (s, 2H) 7.46-7.52 (m, 2H), 7.55 (d, 2H), 7.65-7.73 (m, 4H), 7.80 (d, 1H), 8.76 (m, 1H), 8.85 (d, 1H), 9.12 (s, 1H), 9.82 (br, NH).

EXAMPLE 57

N-Cyclopropyl-1-[3-(1,6-dihydro-6-oxopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




To a solution of N-cyclopropyl-1-[3-(6-benzyloxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 42 in 1,2-dichloroethane (25ml/mmol) was added trifluoroacetic acid (1.5ml/mmol) and the resulting mixture was stirred at 60°C for 18 hours. More trifluoroacetic acid was added (0.75ml/mmol) and heating was continued for a further 24 hours. The cooled mixture was diluted with methylene chloride and saturated aqueous sodium bicarbonate was added, resulting in precipitation of a solid which was filtered. From the filtrate the organic phase was collected and evaporated to a solid which was combined with the previous filtered solid. This mixture was chromatographed on silica gel eluting with 10% methanol in methylene chloride to afford the title compound as a white fluffy solid.

¹H NMR (DMSO-d₆) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.90 (m, 1H), 6.45 (d, 1H), 7.52 (m, 1H), 7.61-7.65 (m, 2H), 7.78 (d, 1H), 7.85 (s, 1H), 7.89-7.93 (m, 2H), 8.74 (d, 1H), 8.78-8.81 (m, 2H), 9.73 (br, NH), other NH >11ppm.

EXAMPLE 58

N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]pyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Step 4 of Example 47, but substituting N-cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 41 for N-cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a white solid.


¹H NMR (DMSO-d₆) δ 0.58 (m, 2H), 0.79 (m, 2H), 2.91 (m, 1H), 5.30 (s, 1H, OH), 7.65 (m, 1H), 7.71-7.79 (m, 3H), 8.12 (d, 1H), 8.23-8.26 (m, 2H), 8.49 (s, 1H), 8.75 (dd, 1H), 8.80 (m, 1H), 8.87 (s, 1H), 8.97 (m, 2H), 9.04 (s, 1H), 9.74 (br, NH).

15

EXAMPLE 59

N-Isopropyl-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 7 for
10 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.


^1H NMR (DMSO- d_6) δ 1.21 (d, 6H), 4.10 (m, 1H), 7.51 (t, 1H), 7.64 (m, 1H), 7.71-7.75 (m, 3H), 7.97 (m, 1H), 8.09 (s, 1H), 8.23 (d, 1H), 8.69-8.77 (m, 3H), 8.84 (s, 1H), 9.66 (br, NH).

15

EXAMPLE 60

N-(2,6-Dichloropyridin-4-yl)-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5




Following the procedure of Step 2 of Example 30, but substituting N-(2,6-dichloropyridin-4-yl)-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 10 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.

^1H NMR (DMSO- d_6) δ 7.51 (m, 1H), 7.69-7.78 (m, 4H), 7.99 (dd, 1H), 8.14 (s, 1H), 8.24 (dd, 1H), 8.70 (s, 1H), 8.73 (s, 2H), 8.84 (m, 2H), 8.99 (s, 1H), 12.05 (br, NH).

EXAMPLE 61

N-Isopropyl-1-[3-(5-carboethoxy-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



10 Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-[3-(5-carboethoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 24 for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the title compound was obtained as a white solid.


¹H NMR (CDCl₃) δ 1.28 (d, 6H), 1.40 (t, 3H), 4.28 (m, 1H), 4.43 (q, 2H), 7.49 (dd, 1H), 7.56 (m, 1H), 7.68 (s, 1H), 7.73 (d, 2H), 8.04 (s, 1H), 8.60 (s, 1H), 8.68 (dd, 1H), 8.77 (s, 1H), 8.82 (d, 1H), 9.01 (s, 1H), 9.61 (br, NH).

15

EXAMPLE 62

N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



10

Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 25 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid


$^1\text{H NMR}$ (CDCl_3) δ 1.29 (d, 6H), 1.60 (s, 6H), 4.11 (brs, 1H), 4.23 (m, 1H), 7.42-7.51 (m, 2H), 7.58 (s, 2H) 7.65 (m, 2H), 8.28 (s, 1H), 8.33 (s, 1H), 8.64 (m, 1H), 8.80 (d, 1H), 8.98 (s, 1H), 9.61 (br, NH).

15

EXAMPLE 63

N-Isopropyl-1-{3-[6-(2-methylpropyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro
[1,8]naphthyridin-4-one-3-carboxamide

5




Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-
{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide
10 from Example 26
for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as an off-
white solid.

¹H NMR (CDCl₃) δ 0.98 (d, 6H), 1.29 (d, 6H), 2.29 (m, 1H), 2.32 (d, 2H), 4.26
(m, 1H), 7.28 (d, 1H), 7.38 (d, 1H), 7.47-7.52 (m, 2H), 7.60 (s, 1H), 7.69 (m, 2H), 8.53 (s, 1H),
15 8.69 (m, 1H), 8.82 (dd, 1H), 9.03 (s, 1H), 9.62 (br, NH).

EXAMPLE 64

N-Isopropyl-1-[3-(6-methyl-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from
10 Example 28 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as an off-white solid.


^1H NMR (CDCl_3) δ 1.32 (d, 6H), 2.60 (s, 3H), 4.30 (m, 1H), 7.35-7.45 (m, 2H), 7.50 (m, 2H), 7.62 (s, 1H), 7.72 (d, 2H), 8.58 (s, 1H), 8.72 (m, 1H), 8.85 (dd, 1H), 9.06 (s, 1H), 9.66 (br, NH).

15

EXAMPLE 65

N-Cyclopropyl-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from
10 Example 14 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.


^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.90 (m, 1H), 7.52 (t, 1H), 7.65 (m, 1H), 7.72-7.76 (m, 3H), 7.98 (m, 1H), 8.10 (s, 1H), 8.25 (d, 1H), 8.70-8.79 (m, 3H), 8.85 (s, 1H), 9.72 (br, NH).

15

EXAMPLE 66

N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Step 2 of Example 29 but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from Step 2 of Example 30 for 5-bromo-1-oxidopyrimidine, the title compound was obtained as a white solid.


10

^1H NMR (CDCl_3) δ 0.66 (m, 2H), 0.85 (m, 2H), 1.70 (s, 6H), 2.97 (m, 1H), 7.43-7.49 (m, 2H), 7.52-7.56 (m, 2H), 7.61 (s, 2H) 7.71-7.74 (m, 2H), 8.49 (s, 1H), 8.68 (m, 1H), 8.80 (d, 1H), 9.02 (s, 1H), 9.74 (br, NH).

EXAMPLE 67

N-Cyclopropyl-1-[3-(1-oxidopyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-
carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-cyclopropyl-1-[3-(pyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from
10 Example 17 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.


^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.79 (m, 2H), 2.92 (m, 1H), 7.62-7.70 (m, 2H), 7.75 (t, 1H), 7.88 (d, 2H), 8.03 (d, 1H), 8.15 (s, 1H), 8.30 (d, 2H), 8.75 (d, 1H), 8.80 (m, 1H), 8.86 (s, 1H), 9.73 (br, NH).

15

EXAMPLE 68

N-Cyclopropyl-1-[3-(5-bromo-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from
10 Example 41 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a light yellow solid.


¹H NMR (DMSO-d₆) δ 0.56 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 7.65 (m, 1H),
7.71-7.74 (m, 2H), 8.02-8.06 (m, 2H), 8.15 (s, 1H), 8.60 (s, 1H), 8.73-8.79 (m, 3H), 8.86 (s,
1H), 9.73 (br, NH).

15

EXAMPLE 69

N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]pyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]pyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 58 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, and using 1.6eq. of m-chloroperoxybenzoic acid, the title compound was obtained as a white solid.


¹H NMR (DMSO-d₆) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 6.94 (s, 1H, OH), 7.65 (m, 1H), 7.71-7.79 (m, 3H), 7.97 (dd, 1H), 8.13 (d, 1H), 8.25 (s, 1H), 8.55 (s, 1H), 8.74 (dd, 1H), 8.80 (m, 1H), 8.87 (s, 1H), 8.91 (s, 1H), 9.00 (s, 1H), 9.09 (s, 1H), 9.73 (br, NH).

15

EXAMPLE 70

N-Cyclopropyl-1-[[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-oxidopyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



From the procedure of Example 69, the title compound was also obtained as a white solid.


10

^1H NMR (DMSO- d_6) δ 0.57 (m, 2H), 0.79 (m, 2H), 2.92 (m, 1H), 5.32 (s, 1H, OH), 7.65 (m, 1H), 7.72-7.80 (m, 3H), 8.08-8.17 (m, 2H), 8.27 (m, 2H), 8.70-8.82 (m, 4H), 8.88 (s, 1H), 8.98 (s, 1H), 9.73 (br, NH).

EXAMPLE 71

N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]-1-oxidopyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



From the procedure of Example 69 the title compound was also obtained as a white solid.


10

^1H NMR (DMSO- d_6) δ 0.58 (m, 2H), 0.80 (m, 2H), 2.92 (m, 1H), 6.85 (brs, 1H, OH), 7.65 (m, 1H), 7.70-7.80 (m, 3H), 7.96 (d, 1H), 8.13 (m, 2H), 8.29 (s, 1H), 8.71-8.84 (m, 4H), 8.89 (s, 1H), 8.92 (s, 1H), 9.73 (br, NH).

EXAMPLE 72

N-Isopropyl-1-[3-(1-oxidoquinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-
carboxamide

5




Following the procedure of Step 2 of Example 30, but substituting N-isopropyl-1-[3-(quinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide from Example 12
10 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a solid.

^1H NMR (CDCl_3) δ 1.30 (d, 6H), 4.28 (m, 1H), 7.49 (dd, 1H), 7.54 (d, 1H), 7.66-7.85 (m, 5H), 7.92 (m, 2H), 8.69-8.75 (m, 2H), 8.84 (d, 1H), 8.86 (s, 1H), 9.08 (s, 1H), 9.64 (br, NH).

EXAMPLE 73

N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide




To a mixture of N-isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 46 in 13:1
 10 methylene chloride/methanol (33ml/mmol) at room temperature was added magnesium
 monoperoxyphthalate hexahydrate (MMPP, 1.1 molareq) and the resulting mixture was stirred at
 room temperature for 24 hours. The mixture was filtered through a bed of celite and the filtrate
 was washed with aqueous sodium carbonate, then water and dried. The crude product was
 chromatographed on silica gel eluting with 8% ethanol in ethyl acetate and the solid obtained
 15 was stirred at room temperature in ether for several hours and filtered to afford the title
 compound as a light pink solid.

¹H NMR (Acetone-d₆) δ 0.98 (d, 6H), 1.61 (s, 6H), 1.88 (m, 1H), 3.26 (t, 2H),
 7.52 (s, 1H, OH), 7.61 (m, 1H), 7.66 (d, 1H), 7.77-7.82 (m, 2H), 7.88 (d, 1H), 7.99 (d, 1H), 8.12
 (s, 1H), 8.68 (s, 1H), 8.73 (m, 1H), 8.80 (dd, 1H), 8.93 (s, 1H), 9.81 (br, NH).

EXAMPLE 74

N-Cyclopropyl-1-[3-(6-methyl-1-oxidopyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Example 73, but substituting N-cyclopropyl-1-[3-(6-methylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 39 for N-isobutyl-1{[3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a solid.


^1H NMR (CDCl_3) δ 0.72 (m, 2H), 0.91 (m, 2H), 2.61 (s, 3H), 3.02 (m, 1H); 7.38 (d, 1H), 7.45 (dd, 1H), 7.49-7.58 (m, 2H), 7.66 (s, 1H), 7.75 (m, 2H), 8.61 (s, 1H), 8.72 (m, 1H), 8.87 (dd, 1H), 9.08 (s, 1H), 9.78 (br, NH).

15

EXAMPLE 75

N-Cyclopropyl-1-[3-(6-methylsulfonyl-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro
[1,8]naphthyridin-4-one-3-carboxamide

5




To a suspension of N-cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 51 in methylene chloride
10 (30ml/mmol) was added urea-hydrogen peroxide (8eq) and the resulting mixture was cooled to 0°C. Trifluoroacetic acid (4.7eq) was added and the mixture was warmed to room temperature as a solution was obtained. After 18 hours, more urea-hydrogen peroxide (2.6eq) and trifluoroacetic acid (2eq) were added and stirring was continued for 2 hours. The mixture was
15 quenched with saturated aqueous sodium metabisulfite, diluted with methylene chloride and the organic phase was washed with 1N aqueous HCl, then brine and water, dried and evaporated. The crude product was chromatographed on silica gel eluting with 40% toluene in acetone to afford the title compound as a solid.

¹H NMR (CDCl₃) δ 0.66 (m, 2H), 0.85 (m, 2H), 2.97 (m, 1H), 3.52 (s, 3H), 7.48 (m, 1H), 7.58-7.65 (m, 3H), 7.72-7.78 (m, 2H), 8.15 (d, 1H), 8.54 (s, 1H), 8.68 (brs, 1H), 8.81
20 (d, 1H), 9.01 (s, 1H), 9.71 (br, NH).

EXAMPLE 76

N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-
dihydro[1,8]naphthyridin-4-one-3-carboxamide

5



Following the procedure of Step 2 of Example 30, but substituting N-cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Example 47 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a solid.


^1H NMR (CDCl_3) δ 0.71 (m, 2H), 0.90 (m, 2H), 1.75 (s, 6H), 3.02 (m, 1H), 7.48-7.60 (m, 5H), 7.73 (s, 1H), 7.88 (s, 1H), 8.52 (s, 1H), 8.72 (m, 1H), 8.84 (dd, 1H), 9.04 (s, 1H), 9.71 (br, NH).

15

EXAMPLE 77

N-Cyclopropyl-1-{3-[6-(1,2-dihydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

5



Step 1: 5-Bromo-2-(1-methylvinyl)pyridine N-oxide

10 A mixture of 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from step 2 of example 30 (1.29g) and 25% aqueous sulfuric acid was heated at 130°C for 2 days. After cooling, the mixture was made slightly basic using 10N aqueous sodium hydroxide and partitioned between ethyl acetate and water. The crude product from evaporation of the organic phase was used as such in step 2.

15 Step 2: 5-bromo-2-(1,2-dihydroxy-1-methylethyl) pyridine N-oxide


The crude product from step 1 was dissolved in a 3 : 1 mixture of acetone and water (16mL) and 4-methylmorpholine N-oxide (1g) and potassium osmate dihydrate (90mg) were added. The resulting mixture was stirred at room temperature for 3 days then excess solid sodium bisulfite was added and the mixture was evaporated. The residue was diluted with 20 methylene chloride and filtered. The filtrate was evaporated and the residue chromatographed on silica gel eluting with ethyl acetate to afford the title compound as a white solid.

Step 3: N-Cyclopropyl-1-{3-[6-(1,2-dihydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

Following the procedure of step 2 of example 32 but substituting 5-bromo-2-(1,2-dihydroxy-1-methylethyl) pyridine N-oxide from step 2 for 3-bromo-5-methylsulfonylpyridine the title compound was obtained as a white solid.


$^1\text{H NMR}$ (CDCl_3) δ 0.66 (m, 2H), 0.85 (m, 2H), 1.61 (s, 3H), 2.78 (m, 1H, OH), 2.97 (m, 1H), 3.90 (m, 1H), 3.97 (m, 1H), 7.48 (m, 1H), 7.53 (m, 2H), 7.60 (m, 2H), 7.69-7.72 (m, 2H), 7.92 (s, 1H, OH), 8.49 (s, 1H), 8.68 (m, 1H), 8.80 (dd, 1H), 9.02 (s, 1H), 9.73 (br, NH).

10 PDE-4 INHIBITOR EXAMPLE B



15 N-Cyclopropyl-1-{3-[6-(1-oxidopyridin-3-yl)ethynyl]phenyl}-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxamide (Compound IX)

Step 1:



	<u>Materials</u>	<u>MW</u>	<u>Amount</u>	<u>Moles</u>	<u>Eq</u>
5	Ethyl ester naphthyridone-phenylacetylene	318.33	2.50 Kg	7.85	1.0
	3-Bromopyridine- <i>N</i> -oxide HCl salt	210.46	1.82 Kg	8.64	1.1
10	<i>Tert</i> -amylamine	87.17 d=0.746	2.7 L	45.6	3.0
15	Allylpalladiumchloride dimer	365.85	72 g	0.2	2.5mol%
	Tri- <i>t</i> -butylphosphine	10% in hexanes (0.33M)	2.38 L	0.8	10mol%
20	Dimethylaminoacetamide solvent (DMAc)		60 L	---	24vol
	water	anti-solvent	90 L	---	36vol
25	n-Butanol	rinse solvent	87.6 L	---	35vol

A 100 L round bottom flask was equipped with a reflux condenser and nitrogen sweep, and charged with bromopyridine-*N*-oxide HCl salt, followed by DMAc 12.5 L, and cooled with ice bath. Neat *t*-amylamine was added via dropping funnel over 1 hour. Addition of amine was noted to be mildly exothermic: temperature rose from 5°C to 10°C. The phosphine solution was added to produce a light yellow, homogeneous solution. Next, the π -allylpalladium dimer was added as a slurry in DMAc 1.5 L. The reaction was allowed to stir at 10°C for 30 minutes. The acetylene compound was weighed out into a carboy and DMAc 9.9 Kg added.

This acetylene solution was added into the reaction flask. The carboy was rinsed with DMAc 22 Kg and added to the reaction mixture. Enough DMAc was added to achieve a total of 60 L. This milky yellow solution was heated to 60°C, becoming a homogeneous brown solution, until reaction was complete, judged by consumption of the acetylene compound. The reaction was cooled to 0°C before adding water 90 L to precipitate product from solution. The slurry was filtered to collect product. To remove palladium, the cake was washed with 87.5 L of n-butanol on the filter pot, and then dried in vacuum oven at 50°C under nitrogen stream overnight to give white solids 2.68 Kg, 83% yield

NMR ¹H (CDCl₃ 400MHz) δ: 1.35 (t, , J=7.1 Hz, 3H), 4.34 (q, J=7.1, 2H), 7.28 (dd, J=8.0, 6.8 Hz, 1H), 7.34 (dT, J=8.0, 1.2 Hz, 1H), 7.38 (dd, J=8.0, 4.4 Hz, 1H), 7.49 (ddd, J=8.0, 2.4, 1.2 Hz, 1H), 7.58 (t, J=7.9 Hz, 1H), 7.64 (t, J=1.8 Hz, 1H), 7.67 (dt, J=7.6, 1.2 Hz, 1H), 8.12 (dt, J=6.5, 1.1 Hz, 1H), 8.27 (s, 1H), 8.59 (dd, J=4.4, 1.9 Hz, 1H), 8.64 (s, 1H), 8.74 (dd, J=8.0, 1.9 Hz, 1H).

NMR ¹³C (CDCl₃ 400MHz) δ: 14.3, 61.1, 84.7, 92.4, 112.8, 121.4, 122.7, 123.0, 123.2, 125.8, 128.3, 128.5, 129.8, 130.7, 132.5, 136.8, 139.1, 141.0, 149.0, 149.9, 152.5, 164.6, 174.5.


An alternative method of Palladium reduction has also been used and is as follows:

<u>Materials</u>	<u>MW</u>	<u>Amount</u>	<u>Moles</u>	<u>Eq</u>
Ester product	411.14	0.718 Kg	7.85	1.0
Darco G-60	---	0.359 Kg	---	50wt%
Methanol	solvent	82 L	---	50vols

After precipitation with water, the wet filter cake was dried under a nitrogen stream. A 100 L round bottom flask was equipped with a reflux condenser, stir paddle, and nitrogen sweep, then charged with the ester product and methanol 72 L. Mixture was heated to 60°C until solids dissolved. Mixture was cooled to 55°C and Darco G-60 was added as a slurry in 6L methanol. Mixture was heated back to 60°C for four hours. The solution was filtered through Solka Floc while warm to remove Darco. Solka Floc pad was rinsed with another 4L of

methanol. Combined filtrates were concentrated to 25L volume, at which point a copious white precipitate was noted. This slurry can be carried forward into the amide formation step directly.

Step 2:



	<u>Materials</u>	<u>MW</u>	<u>Amount</u>	<u>Moles</u>	<u>Eq</u>
10	Penultimate ester	411.41	1.4 Kg	3.4	1.0
	Cyclopropylamine	57.10, d=0.824	21 L	---	15vols
	(Butylphosphite	250, d=0.915	46mL	0.17	5mol%)*
15	Methanol	solvent	21 L	---	15vols
	Isopropyl acetate	anti-solvent	72 L	---	---
20	15% aq. acetonitrile	solvent	24 L	---	24vols

In-line filter, 1 micron

*Note that the parentheses around the butylphosphite data indicate that the experiment was run both with and without the butylphosphite.

25 A 100 L round bottom flask was equipped with a reflux condenser and nitrogen sweep, and charged with penultimate ester and methanol followed by cyclopropylamine. As


indicated above, catalyst or Lewis acid can be added. The reaction was heated to 60°C until consumption of the ester. The reaction was judged to be complete when esters were less than 2% by HPLC. Upon completion, the reaction was allowed to cool to 40°C and concentrated to ~half volume. Isopropyl acetate was added while distilling out the remainder of the methanol and amine. After completion of the solvent switch to isopropyl acetate, the final volume was 20L. The slurry was sampled and filtered. The cake was washed with 2 L of isopropyl acetate and then dried in vacuum oven at 50°C under nitrogen stream overnight.

A 72 L round bottom flask was equipped with a reflux condenser and nitrogen sweep, and charged with Compound IX product solids and 24 L 15% aq. acetonitrile. The Slurry was heated to 60°C. All solids went into solution. Batch solution was filtered through a 1 micron in-line filter while warm into a second 72 L round bottom flask, then allowed to cool to 22°C. The slurry was filtered and solids dried in vacuum oven at 50°C under nitrogen stream overnight to give 0.86 Kg of crystallized Compound IX, 60% yield.

NMR ^1H (CDCl_3 400MHz) δ : 0.61-0.73 (m, 2H), 0.79-0.92 (m, 2H), 2.95-3.01 (m, 1H), 7.28 (dd, $J=8.0, 6.8$ Hz, 1H), 7.40 (d, $J=8.0$ Hz, 1H), 7.48 (dt, $J=8.0, 4.4$ Hz, 1H), 7.49 (ddd, $J=8.0, 2.0, 1.2$ Hz, 1H), 7.61 (t, $J=8.0$ Hz, 1H), 7.64 (m, $J=2.0$ Hz, 1H), 7.71 (dt, $J=7.6, 1.2$ Hz, 1H), 8.19 (d, $J=6.6$, 1H), 8.34 (s, 1H), 8.70 (dd, $J=4.6, 1.8$ Hz, 1H), 8.80 (dd, $J=8.0, 1.9$ Hz, 1H), 9.00 (s, 1H), 9.74 (br d, $J=3.6$ Hz, 1H).

NMR ^{13}C (CDCl_3 400MHz) δ : 6.5, 22.4, 84.6, 92.5, 113.5, 121.4, 122.0, 122.9, 123.2, 125.7, 128.4, 128.6, 129.8, 130.7, 132.6, 136.4, 139.0, 140.4, 141.1, 148.0, 149.7, 153.0, 165.1, 177.1.

Alternative Procedure Illustrating Amide Formation with Magnesium Chloride



	<u>Materials</u>	<u>MW</u>	<u>Amount</u>	<u>Moles</u>	<u>Eq</u>
	Penultimate ester	411.41	1.0 Kg	2.43	1.0
5	Magnesium chloride	95.22	0.231 Kg	2.43	1.0
	Cyclopropylamine	57.10, d=0.824	10 L	---	10vol
	Acetonitrile	solvent	30 L	---	30vol
10			16 L		16vol
	Water	solvent	8 L	---	8vol
	sodium citrate	294.10	0.79 Kg	2.67	1.1
15	In-line filter, 1 micron				

Process Description

20 A 72 L round bottom flask was equipped with a reflux condenser and nitrogen sweep, and charged with penultimate ester and acetonitrile followed by magnesium chloride and then cyclopropylamine. Reaction was heated to 60°C until complete consumption of ester. Reaction was judged complete when esters were less than 0.5% by HPLC. Upon completion, reaction was allowed to cool to 40°C and concentrated to ~half volume. Acetonitrile was added

25 while distilling out remainder of cyclopropylamine. After complete solvent switch to acetonitrile, final volume was 8L. Water 8L was added and slurry was heated to 60°C. All solids went into solution. Batch solution was filtered through a 1 micron in-line filter while warm into a 22 L round bottom flask, then allowed to cool to 22°C. Slurry was filtered and solids dried in vacuum oven at 50°C under nitrogen stream overnight to give 0.82 Kg of '191,


30 80% yield.

Step 3: Recrystallization of Compound IX

	<u>Materials</u>	<u>MW</u>	<u>Amount</u>	<u>Moles</u>	<u>Eq</u>
5	Compound IX	422.45	0.86 Kg	2.0	1.0
	Ethanol	solvent	20 L	---	4.3vols

10 A 72 L round bottom flask was equipped with a reflux condenser and nitrogen sweep, and charged with Compound IX and ethanol. Slurry was heated to 50°C with stirring until recrystallization was complete. Recrystallization was judged to be complete when greater than 95% Form A by solid-state NMR and/or powder diffraction X-ray. The batch was allowed to cool to 22°C, then filtered to collect white solids which were dried in a vacuum oven at 50°C under a nitrogen stream overnight to give 0.74 Kg of the desired form. The solids were co-
 15 milled through a size 12 mesh to provide fluffy white solids 0.719 Kg.

PDE-4 INHIBITOR EXAMPLE 2



20 N-Cyclopropyl-1-{3-[3-(pyridin-3-yl)ethynyl]phenyl}-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxamide (Compound IXa)

25 Compound IXa was prepared by following the procedure as described for Example 1, with the exception that 3-bromopyridine was substituted for 3-bromopyridine-N-oxide.

Other variations or modifications, which will be obvious to those skilled in the art, are within the scope and teachings of this invention. This invention is not to be limited except as set forth in the following claims.